



زانكۆی سه‌لاحه‌دین - هه‌ولێر
Salahaddin University-Erbil

Antibiotic Resistance Pattern and Extended Spectrum Beta Lactamase Producing *Escherichia coli* Isolated From clinical samples

Research Project

Submitted to the Department of (Biology) in partial fulfillment of the requirements for the degree of **BSc. in Biology**

By:

Aran Haidar

Supervised by:

Pishtiwan A. Hamad

April – 2023

CERTIFICATE

This research project has been written under my supervision and has been submitted for the award of the **BSc.** degree in **Biology** with my approval as a supervisor.

Signature

Name:

Date: **April, 2023**

DEDICATION

This effort I dedicate to **Allah** Almighty, my lord, my powerful foundation, my source of inspiration, wisdom, knowledge, and understanding. Throughout this project, he was the source of my energy.

Student Name

Aran Haidar

ACKNOWLEDGMENTS

To begin with, I thank (**Allah**) for His blessing, which made me able to complete and perform this study with success, the lord of the universe, blessing, and peace be on **Muhammad** (Allah's peace and prayers be upon him).

Finally, I want to say thanks to all those I forgot them here to mention his/her name, who assisted me even by one useful scientific word directly or indirectly.

SUMMARY

Background: Due to the recent appearance of organisms that are resistant to several drugs (multidrug-resistant) like Enterobacteriaceae that produce extended-spectrum β -lactamase (ESBL), concerns have remarkably increased regarding the suitable treatment of infections. The present study was an investigation into ESBL molecular characteristics among clinical isolates of *Escherichia coli* and their pattern of antimicrobial resistance in order to come up with helpful information on the epidemiology of these infections and risk factors accompanied with them.

Methods: In order to conduct the study, 10 *E. coli* were isolated and retrieved from Central and Internal Lab. of Teaching Hospital hospitals in Erbil, Iraq during October, 2022 and February, 2023. The collected strains were analyzed, and the profile of their antimicrobial susceptibility was specified. In order to spot β -lactamase genes (i.e., *bla*_{TEM}, *bla*_{SHV}, and *bla*_{CTX-M}), a polymerase chain reaction was conducted.

Results: The findings obtained from multiplex PCR assay showed that out of the collected strains of ESBL-producing *E. coli*, had 100% *bla*_{TEM}, 70% *bla*_{CTX-M} and 60% *bla*_{SHV} genes. Antibiotic resistance pattern of *E. coli* isolates to 10 antibiotics varied widely. However, the majority of the *E. coli* isolates were multi-drug resistant (MDR).

Conclusion: TEM prevalence was high among other types of ESBLs. Overall, the most active antimicrobial agents *in vitro* remained to be the Amikacin.

Keywords: *Escherichia coli*, ESBL, *bla*_{TEM}, *bla*_{SHV}, and *bla*_{CTX-M}

LIST OF CONTENTS

CERTIFICATE	I
DEDICATION	II
ACKNOWLEDGMENTS.....	III
SUMMARY	IV
LIST OF CONTENTS	V
FIGURE.....	VI
TABLE.....	VII
1. INTRODUCTION	1
2. Materials and Methods.....	3
3. Results and Discussion	6
4. CONCLUSIONS.....	12
REFERENCES	13

FIGURE

Figure1 Agarose gel electrophoresis of PCR amplification products for the 16S ribosomal RNA of *E. coli*. M: The DNA marker (100 bp ladder), lanes **1, 2, 3, 4, 5, 6, 7, 8, 9,** and **10** positive amplification of **16S ribosomal RNA**.....7

Figure2 PCR products of SHV, CTX-M and TEM genes with expected sizes of 747 bp, 593 bp and 445 bp respectively against 100 bp DNA Ladder on a 1% agarose gel.
.....9

TABLE

Table1 Antibiotics name, abbreviations, and standard diameter of inhibition zone	4
Table 2 List of primers used for Multiplex PCR amplification.....	5
Table 3 Morphological and biochemical characteristics of <i>E. coli</i> isolates.....	6
Table 4 Percentage Resistance of <i>E. coli</i> to antibiotics.....	8
Table 5 Antibiotic Susceptibility Results of <i>E. coli</i>	9

1. INTRODUCTION

Escherichia coli (*E. coli*) that belong to the Enterobacteriaceae family, was named after the German pediatrician, Theodore Escherich, who first identified and described it in 1885. *E. coli* are Gram-negative, are typically rod-shaped, facultatively anaerobic bacteria (Chakraborty, 2005). *E. coli* is one of the main causes of both nosocomial and community-acquired infections in human (Diekema et al., 1999). However, some specific *E. coli* strains represent primary pathogens with an enhanced potential to cause disease after acquiring specific virulence attributes. These virulence attributes are normally encoded on genetic elements that can be exchanged between different strains (Kaper et al., 2004).

Bacterial resistance to antimicrobial agents is a threat to public health throughout the world. The presence of antibiotic resistance genes on bacterial plasmids and transposons have further helped in the transmission and spread of drug resistance among pathogenic bacteria. During the last years, outbreaks of diseases due to multi-resistant strains of bacterial pathogens have occurred (Reichling et al., 2009).

The discovery of antimicrobial agents had a major impact on the rate of survival from infections. However, the changing patterns of antimicrobial resistance caused a demand for new antibacterial agents. Routine monitoring of antibiotic resistance provides data for antibiotic therapy and resistance control, prescription programs, making policy decisions and assessing the effectiveness of both (Olowe et al., 2008).

Recently, it was found that many strains *E. coli* have the ability to produce plasmid-encoded enzymes called extended spectrum β -lactamases (ESBLs) as one of the most well-known resistance mechanisms in Gram-negative bacilli (Ojdana et al., 2014). ESBLs are a group of enzymes that lead to resistance increase in Aztreonam, Ceftazidime, Cefotaxime, related Oxyimino- β -lactams, cephalosporins, and

penicillins, but Clavulanic acid inhibits them. TEM, SHV, and CTX-M are the 3 main types of ESBLs. CTX-M, which has become more prevalent than SHV and TEM, includes a rapidly expanding family which has spread among a wide range of clinically important bacteria and over wide geographic areas (Paterson and Bonomo, 2005). Furthermore, strains that produce ESBL often demonstrate resistance to antibiotics belonging to other classes (i.e. aminoglycosides, quinolones, and sulfonamides), which makes strategies of treatment more complex (Liao et al., 2017).

In addition, Enterobacteriaceae family members such as *Klebsiella pneumoniae* and *Escherichia coli* often produce ESBLs; however, other genera of the *Enterobacteriaceae* family have recently been reported to contain some other enzymes. A higher level of resistance in such organisms was first observed in patients with prolonged hospital stays in intensive care units in Europe. However, isolates were identified in Africa, Asia, the Middle East, and South and North Americas, and ESBL GNB soon became a global problem and concern (Malloy and Campos, 2011).

Common ESBL genes coding for isolates of *K. pneumoniae* and *E. coli* were determined as CTX-M (cefotaximase that preferentially hydrolyzes cefotaxime), TEM (found and isolated in the early 80s from Teminora who was a Greek patient), and SHV (for variable of sulphhydryl which was first observed in a single *Klebsiella ozaenae* strain retrieved in Germany). These genes which are mediated by transposons, plasmids, or chromosomes are all sporadically described all over the world (Akpaka et al., 2010).

The aims of this study are to isolate and identify *E. coli* from different clinical samples as well as determining the susceptibility pattern and Extended Spectrum Beta Lactamase of the isolates.

2. MATERIALS AND METHODS

Specimen's collection

A total of (10) isolates of *E. coli* were collected from urine and wound, within the period between October, 2022 and February, 2023 in Central and Internal Lab. of Teaching Hospital in Erbil city.

Culture media

Different culture media such as Nutrient agar, MacConkey agar, Eosin methylene blue and Blood agar were used to isolate and identification of this above pathogens, these media were prepared as described by (Atlas, 2010).

Isolation and identification of bacteria

All isolates which collected were cultured on different media (Nutrient agar, MacConkey agar and Blood agar) by streaking method and incubated at 37 °C for 18-24hours. Discrete colonies were sub-culture into nutrient agar aseptically to obtain pure culture of the isolates, then transfer single colonies to MacConkey agar directly and incubated for 24 hours for screening of *E. coli*. Size, shape and fermenting of lactose were studied, and then the colonies that fermenting lactose sugar and carried *E. coli* features were purified more than one time to obtain purified culture. For confined diagnosis of *E. coli* colonies were inoculated on Eosin methylene blue agar with streaking method and incubated at 37C⁰ for 18-24 hours. Then all isolates were identified according to morphology, cultural and biochemical tests. Besides, 16S rRNA-based molecular identification of clinical isolates was utilized to re-identify them.

Susceptibility studies

The isolates were subjected to antibiotic sensitivity test to 10 antibiotics which include The isolates were subjected to antibiotic sensitivity test to 10 antibiotics which include Amikacin (AK 30 µg), Amoxicillin-Clavulanic acid (AMC 20+10 µg), Ceftriaxone (CTR 30 µg), Ciprofloxacin (CIP 5 µg), Gentamycin (G 10 µg), Nitrofurantoin (F 300 µg), Trimethoprim-Sulphamethaxazole (SXT 1.25+23.75 µg), Nalidixic acid (NA 30 µg), Ampicillin (AMP 10 µg), and Tobramycin (TOB 10 µg) as clarified in **Table (1)** was determined according to NCCLS (National Committee for Clinical Laboratory

Standards) standards. Twenty ml of Muller Hinton agar melted and cooled at 45 °C was poured into sterile petri plates and allowed to solidify completely. A lawn of test pathogen was prepared by evenly spreading 100µl inoculums (1.5 x 10⁸ CFU/ml) (according to 0.5 McFarland standard solutions (McFarland, 1907) with the help of a sterilized swab onto the entire surface of the agar plate. The plates were allowed to dry before applying antibiotic disc. The discs were firmly applied to the surface of agar plates within 15 minutes of inoculation(Wayne, 2012).

Table (1): Antibiotics name, abbreviations, and standard diameter of inhibition zone

No.	Antibiotics	Symbol	Disk potency (µg or U)	Zone Diameter		
				Resistant	Intermediate	Sensitive
				(mm)	(mm)	(mm)
1	Amikacin	AK	30	≤ 14	15 – 16	≥ 17
2	Amoxicillin - Clavulanic acid	AMC	20+10	≤ 13	14 – 19	≥ 20
3	Ceftriaxone	CTR	30	≤ 15	16 – 20	≥ 21
4	Ciprofloxacin	CIP	5	≤ 21	22 – 24	≥ 25
5	Gentamycin	G	10	≤ 15		≥ 15
6	Nitrofurantoin	F	300	≤ 14	15 – 16	≥ 17
7	Trimethoprim – Sulphamethaxazole	SXT	1.25+23.75	≤ 10	11 – 15	≥ 16
8	Nalidixic acid	NA	30	≤ 25		≥ 26
9	Ampicillin	AMP	10			
10	Tobramycin	TOB	10	≤ 12	13-14	≥ 15

Amplification of 16S gene

Universal primers 1492R (1) (5'- GGTTACCTTGTTACGACTT -3') as reverse primer and 8F (5'- AGAGTTTGATCCTGGCTCAG -3') as forward were used to carry out PCR for the amplification of the 16S rRNA gene (Turner et al., 1999). All reactions of PCR were conducted by utilizing 2 µl DNA template (density of 10 ng/µl), the Master Mix consisting of 3 mM MgCl₂, 0.2% Tween® 20, 20 mM Tris-HCl pH 8.5, (NH₄)₂SO₄, 0.2 units/µl Ampliqon Taq DNA polymerase, 0.4 µM of each primer, and 0.4 mM of each dNTP. The conditions of PCR included primary denaturation at 95 °C for 5 min, followed by 35 cycles at 95 °C for 45 s, at 55 °C for 45 s, and at 72 °C for 90 s, and a last extension at 72 °C for 6 min. After safe staining, the product of the reaction was

visualized under UV light on a 1% agarose gel (Russell and Sambrook, 2001) (**Fig. 1**). The PCR products were used for sequencing the 16S rRNA of all 78 bacterial isolates.

Detection of ESBL genotypes by multiplex PCR amplification

Using the method utilized by (Monstein et al., 2007) with slight modifications, multiplex PCR was employed to examine the positive isolates in the initial screening test for ESBL production for the existence of *bla*SHV, *bla*CTX-M, and *bla*TEM genes. Freshly cultured isolates bacteria were used to prepare template deoxyribonucleic acid (DNA) was prepared using Presto™ Mini gDNA bacterial kit. All reactions of PCR were conducted by utilizing 2 µl DNA template (density of 10 ng/µl), the Master Mix consisting of 3 mM MgCl₂, 0.2% Tween® 20, 20 mM Tris-HCl pH 8.5, (NH₄)₂SO₄, 0.4 mM of each dNTP, 0.4 µM of each primer, and 0.2 units/µl Ampliqon Taq DNA polymerase. The conditions of polymerase chain reaction amplification were set up as follow: primary denaturation step for 10 minutes at 95° C; 30 denaturation cycles for 30 seconds at 94° C, annealing 30 seconds at 60° C for, extension for 2 minutes at 72° C, and a final extension step for 10 minutes at 72° C. Using agarose gel electrophoresis, size separation PCR amplicons were utilized to detect respective genes (**Table2**).

Table 2. List of primers used for Multiplex PCR amplification

Target gene	Primer	Sequence (5'-3')	Amplicon size	References
<i>bla</i> TEM	Forward	TCG CCG CAT ACA CTA TTC TCA GAA TGA	445-bp	(Al-Ouqaili, 2018)
	Reverse	ACG CTC ACC GGC TCC AGA TTT AT		
<i>bla</i> SHV	Forward	ATG CGT TATATT CGC CTG TG	747-bp	(Paterson et al., 2003)
	Reverse	TGC TTT GTT ATT CGG GCC AA		
<i>bla</i> CTX-M	Forward	ATG TGC AGY ACC AGT AAR GTK ATG GC	593-bp	(Al-Ouqaili, 2018)
	Reverse	TGG GTR AAR TAR GTS ACC AGA AYC AGC GG		

3. RESULTS AND DISCUSSION

Ten isolates of *E. coli* were collected from urine and wound of human infections, from Central and Internal Lab. of Teaching Hospital. The isolates were initially been identified using conventional (which is depend on cultural, morphological, and biochemical tests). The characteristics of isolates were studied, through culturing it on MacConkey agar the results showed that the colonies of *E. coli* were circular, lactose fermenter, and rose-pink in color, on EMB agar the colonies were small, flat with a metallic green sheen, as shown by (Brooks et al., 2007). Also, it is cleared from the Gram stain procedure that the bacterial cell of *E. coli* from smear preparation are gram-negative rods as described in (Brooks et al., 2007).

After isolates had initially been identified using conventional, to emphasize the results of isolates under study the molecular method was used for isolate identification based on the sequence of 16S ribosomal RNA. The 16S rRNA gene was amplified for each strain to confidently identify bacteria, figure 1. And the results were recorded in Table (3).

Table3: Morphological and biochemical characteristics of *E. coli* isolates.

Parameters	Results
Gram stain	Negative
Cellular morphology	Rods
Growth on Blood agar	Large grayish-white
Growth on MacConkey agar	Rose-pink
Indole test	Positive
Citrate test	Negative
Urease test	Negative
Catalase test	Positive




Figure 1 Agarose gel electrophoresis of PCR amplification products for the 16S ribosomal RNA of *E. coli*. **M**: The DNA marker (100 bp ladder), lanes **1, 2, 3, 4, 5, 6, 7, 8, 9, and 10** positive amplification of **16S ribosomal RNA**.

Antibiotic susceptibility assay

In the present study table (4) shows the susceptibilities of 10 isolates to different antibiotics and the result showed that all *E. coli* isolates were sensitive to Amikacin and 80% of isolates were sensitive to Gentamicin, then Ceftriaxone Ciprofloxacin by 60%, followed by Tobramycin in which 50% of isolates were sensitive to antibiotics and 40% of isolates were sensitive to Amoxicillin-Clavulanic acid, Nitrofurantoin, Trimethoprim–Sulphamethaxazole and Nalidixic. On the other hand, this study revealed that all the isolates (100%) were resistant to Ampicillin. The resistance of *E. coli* to different antibiotics in this study confirms similar findings reported by several workers. A study done by (Khadri and Alzohairy, 2009) reported that 88%, 72%, 56%, 50%, 46%, 38%, 30%, and 20% of their *E. coli* isolates were sensitive to AK, G, CIP, TOB, CTR, F, SXT, and AMP respectively. Also (Habrun et al., 2010) found in their study that 80% of the isolates were resistance to NA. And finally, a study by (Kaftandzhieva et al., 2009) found that 76% of the isolates were sensitive to AMC.

The high resistance of the bacteria to different antibiotics may be related to the presence and dissemination of plasmids within the heterogeneous population of these bacteria. Our findings agree with other observations done by (Sharma et al., 2007) who demonstrated that using inaccurate concentration of antibiotics or drug or unnecessary of medicine appointment leads to the resistance of sensitive bacteria, in addition to weakling immunity system in some human due to poor nutrition or heredity factors make bacteria to be more resist. On the other hand, the results may regard the increasing resistance of *E. coli* to antibiotics due to the evolution and spread of plasmid-encoded ES β L enzyme and other genes conferring cross-resistance to other antibiotics. This is of concern due to the increasing cost of antibiotic treatment and the spread of multi-drug resistance to more pathogenic microorganisms (Kamatchi et al., 2009). ES β L is enzymes which can hydrolyze Penicillins as well as Cephalosporins and Monobactam. The majority of ES β L isolates are also resistant to other antibiotics such as Fluoroquinolones and Aminoglycosides. This finding was supported by those previously recorded by (Chaudhary and Aggarwal, 2004).

Also, the table (5) shows a detailed resistance pattern to antimicrobial agents. All isolates of *E. coli* show difference in susceptibility against these antibiotics which used in this study started from 10% of resistant as a low percentage and a high percentage of resistance was recorded by isolate no. 2 which represent 80% to all antibiotics.

Table (4): Percentage Resistance of *E. coli* to antibiotics

No.	Antibiotics*	Sensitive (%)	Resistant (%)
1	AK	10(100)	0(00)
2	AMC	4(40)	6(60)
3	CTR	6(60)	4(40)
4	CIP	6(60)	4(40)
5	G	8(80)	2(20)
6	F	4(40)	6(60)
7	SXT	4(40)	6(60)
8	NA	4(40)	6(60)
9	AMP	0(00)	10(100)
10	TOB	5(50)	5(50)

Table (5): Antibiotic Susceptibility Results of *E. coli*

Bacterial isolate no.	Antibiotics										% Resistance
	AK	AMC	CTR	CIP	G	F	SXT	NA	AMP	TOB	
ISO 1	S	S	S	S	R	S	S	s	R	S	20
ISO 2	S	R	S	R	R	R	R	R	R	R	80
ISO 3	S	R	R	S	S	R	R	R	R	S	60
ISO 4	S	S	S	S	S	S	S	S	R	S	10
ISO 5	S	R	R	S	S	R	S	R	R	R	60
ISO 6	S	R	S	R	S	R	R	R	R	S	60
ISO 7	S	S	S	S	S	S	S	S	R	R	20
ISO 8	S	R	R	R	S	R	R	S	R	S	60
ISO 9	S	R	S	R	S	S	R	R	R	R	60
ISO 10	S	S	R	S	S	R	R	R	R	R	60
% Sensitive	100	40	60	60	80	40	40	40	0	50	

In PCR detection of ESBL genotypes, it was found that all of the ESBL screening positive *E. coli* isolates had one or more ESBL genes that were tested in the present study. Overall, 100% (10/10) of *E. coli* isolates were positive for one or more ESBL genes. The multiplex PCR assay results indicated that 100% *bla*TEM genes, 70% *bla*CTX-M, and 60% *bla*SHV genes were detected in the *E. coli* isolates. The overall incidence of ESBL genotypes in *E. coli* isolates is illustrated in **figure 2**.




Figure 2. PCR products of SHV, CTX-M and TEM genes with expected sizes of 747 bp, 593 bp and 445 bp respectively against 100 bp DNA Ladder on a 1% agarose gel.

Phenotypic tests for detection of ESBL can only confirm ESBL production but fail to recognize the subtypes of ESBL. As reported by (Nüesch-Inderbinen et al., 1996), molecular methods have been proved to be sensitive, but they costly and conducting them requires a long time, expertise, and specialized equipment. Ultimate identification is only probable through methods of molecular detection. The results of a study conducted by (Navon-Venezia et al., 2003) revealed that it is necessary to periodically evaluate these phenotypic tests because introduction of new enzyme can change their performance. In their study of phenotypic and genotypic methods of ESBL detection, (Grover et al., 2006) stated that PCR is a reliable method for detecting ESBL. In the present study, multiplex PCR amplification assay was utilized to detect blaCTX-M, blaSHV, and blaTEM genes in the retrieved clinical isolates of *E. coli* because one of the advantages of this assay rapid screening of large numbers of clinical isolates, moreover, if it is required, further molecular epidemiological studies can take advantage of the DNA that is isolated via this assay (Monstein et al., 2007).

Furthermore, it is essential to identify beta lactamase in order to conduct a reliable epidemiological investigation into antimicrobial resistance. The current study was conducted to survey antimicrobial drug resistance, ESBL phenotypes, and blaSHV, blaTEM and blaCTX-M genes detection in *E. coli* isolates retrieved from urinary tract infections in Erbil.

The most globally common type of ESBL appeared to be CTX-M-type ESBLs with their higher incidence in most locations compared to SHV and TEM ESBLs (Jorgensen et al., 2010). Among the three ESBL genotypes included in this study, the most prevalent one was found to be blaTEM (100%) in ESBL-producing isolates of *E. coli*. The less prevalent ESBL genotype was blaSHV, and the prevalence rate of blaSHV in ESBL-producing *E. coli* isolates (60%). A study conducted by (Manoharan et al., 2011) reported similar findings. Furthermore, in another study, (Moghnieh et al., 2018) have reported that *E. coli* and *Klebsiella* spp resistance to third-generation

cephalosporins is usual in whole countries, with outbreak reaching over 50% in Egypt and Syria.

4. CONCLUSIONS

Knowledge of the antimicrobial resistance patterns and resistance genes of bacterial pathogens in a geographical area is important for control and surveillance of antibiotic resistance. The results of the present study revealed that MDR was highly prevalent. In addition, the Carbapenems, Amikacin, and Ciprofloxacin were found to be the most to the least active antimicrobial agents in vitro. Based on the results obtained in the present study, TEM was highly prevalent among other types of ESBLs.

5. REFERENCES

Akpaka, P., Legall, B., and Padman, J. (2010). Molecular detection and epidemiology of extended-spectrum *beta*-lactamase genes prevalent in clinical isolates of *Klebsiella pneumoniae* and *E coli* from Trinidad and Tobago. *West Indian Medical Journal* **59**, 591-596.

Al-Ouqaili, M. T. (2018). Molecular detection and sequencing of SHV gene encoding for extended-spectrum β -lactamases produced by multidrug resistance some of the Gram-negative bacteria. *International Journal of Green Pharmacy (IJGP)* **12**.

Atlas, R. M. (2010). "Handbook of microbiological media," CRC press.

Brooks, G., Carroll, K., Butel, J., Morse, S., and Mietzner, T. (2007). Jawetz, Melnick, & Adelberg's Medical Microbiology, The McGraw-Hill Companies. *Inc., USA*.

Chakraborty, P. (2005). "A textbook of Microbiology," New Central Book Agency.

Chaudhary, U., and Aggarwal, R. (2004). Extended spectrum β -lactamases (ESBL)–an emerging threat to clinical therapeutics. *Indian journal of medical microbiology* **22**, 75-80.

Diekema, D., Pfaller, M., Jones, R., Doern, G., Winokur, P., Gales, A. C., Sader, H. S., Kugler, K., Beach, M., and Group, S. P. (1999). Survey of bloodstream infections due to gram-negative bacilli: frequency of occurrence and antimicrobial susceptibility of isolates collected in the United States, Canada, and Latin America for the SENTRY Antimicrobial Surveillance Program, 1997. *Clinical Infectious Diseases* **29**, 595-607.

Grover, S., Sharma, M., Chattopadhyaya, D., Kapoor, H., Pasha, S., and Singh, G. (2006). Phenotypic and genotypic detection of ESBL mediated cephalosporin resistance in *Klebsiella pneumoniae*: emergence of high resistance against cefepime, the fourth generation cephalosporin. *Journal of Infection* **53**, 279-288.

- Habrun, B., Kompes, G., Cvetnić, Ž., Špičić, S., BeniĆ, M., and Mitak, M. (2010). Antimicrobial sensitivity of *Escherichia coli*, *Salmonella* spp., *Pasteurella multocida*, *Streptococcus suis* and *Actinobacillus pleuropneumoniae* isolated from diagnostic samples from large pig breeding farms in Croatia. *Veterinarski arhiv* **80**, 571-583.
- Jorgensen, J. H., McElmeel, M., Fulcher, L., and Zimmer, B. (2010). Detection of CTX-M-type extended-spectrum beta-lactamase (ESBLs) by testing with MicroScan overnight and ESBL confirmation panels. *Journal of Clinical Microbiology* **48**, 120-123.
- Kaftandzhieva, A., Kotevska, V., Jankoska, G., Kjurcic-Trajkovska, B., Cekovska, Z., and Petrovska, M. (2009). Extended-spectrum beta-lactamase-producing *E. coli* and *Klebsiella pneumoniae* in children at University pediatric clinic in Skopje. *Macedonian Journal of Medical Sciences*.
- Kamatchi, C., Magesh, H., and Sekhar, U. (2009). Identification of clonal clusters of *Klebsiella pneumoniae* isolates from Chennai by extended spectrum beta lactamase genotyping and antibiotic resistance phenotyping analysis.
- Kaper, J. B., Nataro, J. P., and Mobley, H. L. (2004). Pathogenic *Escherichia coli*. *Nature reviews microbiology* **2**, 123.
- Khadri, H., and Alzohairy, M. (2009). High prevalence of multi-drug-resistance (MDR) and extended spectrum β -lactamases (ESBL) producing bacteria among communityacquired urinary tract infections (CAUTI). *Journal of bacteriology Research* **1**, 105-110.
- Liao, K., Chen, Y., Wang, M., Guo, P., Yang, Q., Ni, Y., Yu, Y., Hu, B., Sun, Z., and Huang, W. (2017). Molecular characteristics of extended-spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* causing intra-abdominal infections from 9 tertiary hospitals in China. *Diagnostic Microbiology and Infectious Disease* **87**, 45-48.

- Malloy, A. M., and Campos, J. M. (2011). Extended-spectrum beta-lactamases: a brief clinical update. *The Pediatric Infectious Disease Journal* **30**, 1092-1093.
- Manoharan, A., Premalatha, K., Chatterjee, S., Mathai, D., and Group, S. S. (2011). Correlation of TEM, SHV and CTX-M extended-spectrum beta lactamases among Enterobacteriaceae with their in vitro antimicrobial susceptibility. *Indian J Med Microbiol* **29**, 161-4.
- Moghnieh, R. A., Kanafani, Z. A., Tabaja, H. Z., Sharara, S. L., Awad, L. S., and Kanj, S. S. (2018). Epidemiology of common resistant bacterial pathogens in the countries of the Arab League. *The lancet infectious diseases* **18**, e379-e394.
- Monstein, H. J., Östholm-Balkhed, Å., Nilsson, M., Nilsson, M., Dornbusch, K., and Nilsson, L. (2007). Multiplex PCR amplification assay for the detection of *bla*SHV, *bla*TEM and *bla*CTX-M genes in Enterobacteriaceae. *Apmis* **115**, 1400-1408.
- Navon-Venezia, S., Hammer-Munz, O., Schwartz, D., Turner, D., Kuzmenko, B., and Carmeli, Y. (2003). Occurrence and phenotypic characteristics of extended-spectrum β -lactamases among members of the family Enterobacteriaceae at the Tel-Aviv Medical Center (Israel) and evaluation of diagnostic tests. *Journal of Clinical Microbiology* **41**, 155-158.
- Nüesch-Inderbinnen, M., Hächler, H., and Kayser, F. (1996). Detection of genes coding for extended-spectrum SHV beta-lactamases in clinical isolates by a molecular genetic method, and comparison with the E test. *European Journal of Clinical Microbiology and Infectious Diseases* **15**, 398-402.
- Ojdana, D., Sacha, P., Wiczorek, P., Czaban, S., Michalska, A., Jaworowska, J., Jurczak, A., Poniatowski, B., and Trynieszewska, E. (2014). The occurrence of *bla*CTX-M, *bla*SHV, and *bla*TEM genes in extended-spectrum β -lactamase-positive strains of *Klebsiella pneumoniae*, *Escherichia coli*, and *Proteus mirabilis* in Poland. *International Journal of Antibiotics* **2014**.

- Olowe, O., Okanlawon, B., Olowe, R., and Olayemi, A. (2008). Antimicrobial resistant pattern of *Escherichia coli* from human clinical samples in Osogbo, south western Nigeria. *African Journal of Microbiology Research* **2**, 8-11.
- Paterson, D. L., and Bonomo, R. A. (2005). Extended-spectrum β -lactamases: a clinical update. *Clinical Microbiology Reviews* **18**, 657-686.
- Paterson, D. L., Hujer, K. M., Hujer, A. M., Yeiser, B., Bonomo, M. D., Rice, L. B., and Bonomo, R. A. (2003). the International *Klebsiella* Study Group: Extended-spectrum β -lactamases in *Klebsiella pneumoniae* bloodstream isolates from seven countries: Dominance and widespread prevalence of SHV-and CTX-M-type β -lactamases. *Antimicrob Agents Chemother* **47**, 3554-3560.
- Reichling, J., Schnitzler, P., Suschke, U., and Saller, R. (2009). Essential oils of aromatic plants with antibacterial, antifungal, antiviral, and cytotoxic properties—an overview. *Complementary Medicine Research* **16**, 79-90.
- Russell, D. W., and Sambrook, J. (2001). "Molecular cloning: a laboratory manual," Cold Spring Harbor Laboratory Cold Spring Harbor, NY.
- Sharma, S., Bhat, G., and Shenoy, S. (2007). Virulence factors and drug resistance in *Escherichia coli* isolated from extraintestinal infections. *Indian journal of medical microbiology* **25**, 369-373.
- Turner, S., Pryer, K. M., Miao, V. P., and Palmer, J. D. (1999). Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis 1. *Journal of Eukaryotic Microbiology* **46**, 327-338.
- Wayne, P. (2012). USA: Clinical and Laboratory Standards Institute (CLSI). 2012. *M100-S21: Performance standards for antimicrobial susceptibility testing. 21st Informational Supplement*.