

# On Soft $p_c$ -Regular and Soft $p_c$ -Normal Spaces

Qumri H. Hamko<sup>†</sup>, Nehmat K. Ahmed<sup>†</sup> and Alias B. Khalaf<sup>‡,1</sup>

<sup>†</sup>Department of Mathematics, College of Education,  
Salahaddin University, Kurdistan-Region, Iraq  
qumri.hamko@su.edu.krd,    nehmat.ahmed@su.edu.krd

<sup>‡</sup>Department of Mathematics, College of Science,  
University of Duhok, Kurdistan Region-Iraq  
aliasbkhalaf@uod.ac

**Abstract :** The aim of this paper is to introduce two new types of soft separation axioms called soft  $p_c$ -regular and soft  $p_c$ -normal spaces by using the concept of soft  $p_c$ -open sets in soft topological spaces. We explore several properties and relations of such spaces. Also we investigate hereditary and soft invariance properties by considering certain soft mappings.

**Keywords :** soft  $p_c$ -open set; soft  $p_c$ -regular space; soft  $p_c$ -normal space.

**2010 Mathematics Subject Classification :** Primary 54A05, 54A10; Secondary 54C05

---

## 1 Introduction

Molodtsov [21] initiated the concept of soft set theory in 1999 as a new mathematical tool to treat many complicated problems related to probability and fuzzy set theory. After that many researchers presented applications of soft set theory in many fields of mathematics such as operations researches, mathematical analysis and algebraic structures. Shabir and Naz [24] in 2011 applied the notion of soft sets to introduce the concept of soft topological spaces which are defined over an initial universe with a fixed set of parameters. They introduced almost all the essential classical notions in topology and defined the concept of soft open sets, soft closed sets, soft interior point, soft closure and soft separation axioms. Al-shami et al [6, ?] investigated several types of soft separation axioms and studied their images and pre-images under soft mappings.

Husain and Ahmed [16] in 2015 studied the properties of soft interior, soft closure and soft boundary operators and they introduced separation axioms by using

---

<sup>1</sup>Corresponding author email: aliasbkhalaf@uod.ac (Alias B. Khalaf)

ordinary points in the universal set also Georgiou et. al [12] in 2013, studied some soft separation axioms, soft continuity in soft topological spaces using ordinary points of a topological space  $X$ . Bayramov et al in [10], defined the notion of soft points and applied them to discuss the properties of soft interior, soft closure and soft boundary operators. They also defined and introduced soft neighborhoods and soft continuity in soft topological spaces using soft points.

It is noticed that a soft topological space gives a parametrized family of topologies on the initial universe but the converse is not true i.e. if some topologies are given for each parameter, we cannot construct a soft topological space from the given topologies. Consequently we can say that the soft topological spaces are more generalized than the classical topological spaces for more details we refer to [5] and [7].

Recently, Hamko and Ahmed [1] introduced the notion of soft  $p_c$ -open sets. They applied this notion to define and discuss the concept of soft  $p_c$ -interior, soft  $p_c$ -closure and soft  $p_c$ -boundary operators. Also they introduced the concept of soft continuity and almost soft continuity by employing soft points and soft  $p_c$ -open sets in a soft topological space.

The aim of this paper, is to introduce and discuss a study of soft separation axioms which we call them soft  $p_c$ -regular and soft  $p_c$ -normal spaces which are defined over an initial universe with a fixed set of parameters. We indicate the relationships between them and present several of their properties.

Throughout the present paper,  $X$  will be a nonempty initial universal set and  $E$  will be a set of parameters. A pair  $(F, E)$  is called a soft set over  $X$ , where  $F$  is a mapping  $F : E \rightarrow P(X)$ . The collection of soft sets  $(F, E)$  over a universal set  $X$  with a parameter set  $E$  is denoted by  $SP(X)_E$ . Any logical operation  $(\lambda)$  on soft sets in soft topological spaces are denoted by usual set theoretical operations with symbol  $(\tilde{s}(\lambda))$ .

## 2 Preliminaries

In this section we present the main definitions and results which will be used in the sequel. For some definitions or results which are not mentioned in this section, we refer to [2], [5, 6], [10], [15], [20] and [25].

**Definition 2.1.** [24] A soft set  $(F, E)$  over  $X$  is said to be null soft set denoted by  $\tilde{\phi}$  if for all  $e \in E$ ,  $F(e) = \phi$  and  $(F, E)$  over  $X$  is said to be absolute soft set denoted by  $\tilde{X}$  if for all  $e \in E$ ,  $F(e) = X$ .

**Definition 2.2.** [24] The complement of a soft set  $(F, E)$  is denoted by  $(F, E)^c$  or  $\tilde{X} \setminus (F, E)$  and is defined by  $(F, E)^c = (F^c, E)$  where  $F^c : E \rightarrow P(X)$  is a mapping given by  $F^c(e) = X \setminus F(e)$ , for all  $e \in E$ .

It is clear that  $((F, E)^c)^c = (F, E)$ ,  $\tilde{\phi}^c = \tilde{X}$  and  $\tilde{X}^c = \tilde{\phi}$ .

**Definition 2.3.** [24] For two soft sets  $(F, E)$  and  $(G, B)$  over a common universe  $X$ , we say that  $(F, E)$  is a soft subset of  $(G, B)$ , if

1.  $E \subseteq B$  and
2. for all  $e \in E$ ,  $F(e) \subseteq G(e)$

We write  $(F, E) \sqsubseteq (G, B)$ .

**Definition 2.4.** [24] The union of two soft sets of  $(F, E)$  and  $(G, B)$  over the common universe  $X$  is the soft set  $(H, C) = (F, E) \sqcup (G, B)$ , where  $C = E \cup B$  and for all  $e \in C$ ,

$$H(e) = \begin{cases} F(e) & : \text{if } e \in E - B \\ G(e) & : \text{if } e \in B - E \\ F(e) \cup G(e) & : \text{if } e \in E \cap B \end{cases}$$

In particular,  $(F, E) \sqcup (G, E) = F(e) \cup G(e)$  for all  $e \in E$ .

**Definition 2.5.** [24] The intersection  $(H, C)$  of two soft sets  $(F, E)$  and  $(G, B)$  over a common universe  $X$ , denoted  $(F, E) \cap (G, B)$ , is defined as  $C = E \cap B$ , and  $H(e) = F(e) \cap G(e)$  for all  $e \in C$ .

In particular,  $(F, E) \cap (G, E) = F(e) \cap G(e)$  for all  $e \in E$ .

**Definition 2.6.** [10] Let  $x \in X$ , then  $(x, E)$  denotes the soft set over  $X$  for which  $x(e) = \{x\}$ , for all  $e \in E$ .

Let  $(F, E)$  be a soft set over  $X$  and  $x \in X$ . We say that  $x \tilde{\in} (F, E)$  read as  $x$  belongs to the soft set  $(F, E)$  whenever  $x \in F(e)$  for all  $e \in E$ .

**Definition 2.7.** [10] The soft set  $(F, E)$  is called a soft point, denoted by  $(x_e, E)$  or  $x_e$ , if for the element  $e \in E$ ,  $F(e) = \{x\}$  and  $F(e) = \phi$  for all  $e \in E \setminus \{e\}$ . We say that  $x_e \tilde{\in} (G, E)$  if  $x \in G(e)$ .

Two soft points  $x_e$  and  $y_{e'}$  are distinct if either  $x \neq y$  or  $e \neq e'$ .

**Remark 2.8.** From Definition 2.6 and Definition 2.7, it is clear that:

1.  $(x, E)$  is the smallest soft set containing  $x$ .
2. if  $x \tilde{\in} (F, E)$  then  $x_e \tilde{\in} (F, E)$ .
3.  $(F, E) = \sqcup \{(x_e, E) : e \in E\}$ .

**Definition 2.9.** [24] Let  $\tilde{\tau}$  be a collection of soft sets over a universe  $X$  with a fixed set  $E$  of parameters. Then  $\tilde{\tau} \subseteq SP(X)_E$  is called a soft topology if,

1.  $\tilde{\phi}$  and  $\tilde{X}$  belongs to  $\tilde{\tau}$ .

2. The union of any number of soft sets in  $\tilde{\tau}$  belongs to  $\tilde{\tau}$ .
3. The intersection of any two soft sets in  $\tilde{\tau}$  belongs to  $\tilde{\tau}$ .

The triplet  $(X, \tilde{\tau}, E)$  is called a soft topological space over  $X$ . The members of  $\tilde{\tau}$  are called soft open sets in  $\tilde{X}$  and complements of them are called soft closed sets in  $\tilde{X}$  and they are denoted by  $SO(\tilde{X})$  and  $SC(\tilde{X})$  respectively. Soft interior and soft closure are denoted by  $\tilde{sint}$  and  $\tilde{scl}$  respectively.

**Definition 2.10.** [24] Let  $(X, \tilde{\tau}, E)$  be a soft topological space and let  $(G, E)$  be a soft set. Then

1. The soft closure of  $(G, E)$  is the soft set  

$$\tilde{scl}(G, E) = \sqcap \{(K, B) \in SC(\tilde{X}) : (G, E) \sqsubseteq (K, B)\}$$
2. The soft interior of  $(G, E)$  is the soft set  

$$\tilde{sint}(G, E) = \sqcup \{(H, B) \in SO(\tilde{X}) : (H, B) \sqsubseteq (G, E)\}.$$

**Definition 2.11.** [15] Let  $(X, \tilde{\tau}, E)$  be a soft topological space,  $(G, E)$  be a soft set over  $\tilde{X}$  and  $x_e \in \tilde{X}$ . Then  $(G, E)$  is said to be a soft neighborhood of  $x_e$  if there exists a soft open set  $(H, E)$  such that  $x_e \in (H, E) \sqsubseteq ((G, E))$ .

**Proposition 2.1.** [24] Let  $(Y, \tilde{\tau}_Y, E)$  be a soft subspace of a soft topological space  $(X, \tilde{\tau}, E)$  and  $(F, E) \in SP(X)_E$ . Then:

1. If  $(F, E)$  is a soft open set in  $\tilde{Y}$  and  $\tilde{Y} \in \tilde{\tau}$ , then  $(F, E) \in \tilde{\tau}$ .
2.  $(F, E)$  is a soft open set in  $\tilde{Y}$  if and only if  $(F, E) = \tilde{Y} \sqcap (G, E)$  for some  $(G, E) \in \tilde{\tau}$ .
3.  $(F, E)$  is a soft closed set in  $\tilde{Y}$  if and only if  $(F, E) = \tilde{Y} \sqcap (H, E)$  for some soft closed  $(H, E)$  in  $\tilde{X}$ .

**Definition 2.12.** [17] A soft subset  $(F, E)$  of a soft space  $\tilde{X}$  is said to be soft pre-open if  $(F, E) \sqsubseteq \tilde{sint}(\tilde{scl}(F, E))$ . The complement of a soft pre-open set is said to be soft pre-closed. The family of soft pre-open (soft pre-closed) set is denoted by  $\tilde{sPO}(X)$  ( $\tilde{sPC}(X)$ ).

**Lemma 2.13.** [17] Arbitrary union of soft pre-open sets is a soft pre-open set.

**Lemma 2.14.** [2] A subset  $(F, E)$  of a soft topological spaces  $(X, \tilde{\tau}, E)$  is soft pre-open if and only if there exists a soft open set  $(G, E)$  such that  $(F, E) \sqsubseteq (G, E) \sqsubseteq \tilde{scl}(F, E)$ .

**Lemma 2.15.** [2] Let  $(F, E) \sqsubseteq \tilde{Y} \sqsubseteq \tilde{X}$ , where  $(X, \tilde{\tau}, E)$  is a soft topological space and  $\tilde{Y}$  is a soft pre-open subspace of  $\tilde{X}$ . Then  $(F, E) \in \tilde{sPO}(X)$ , if and only if  $(F, E) \in \tilde{sPO}(Y)$ .

**Theorem 2.16.** [22] If  $(U, E)$  is soft open and  $(F, E)$  is soft pre-open in  $(X, \tilde{\tau}, E)$ , then  $(U, E) \sqcap (F, E)$  is soft pre-open.

**Lemma 2.17.** [1] Let  $(F, E) \sqsubseteq \tilde{Y} \sqsubseteq \tilde{X}$ , where  $(X, \tilde{\tau}, E)$  is a soft topological space and  $\tilde{Y}$  is a soft subspace of  $\tilde{X}$ . If  $(F, E) \tilde{\in} \tilde{s}PO(X)$ , then  $(F, E) \tilde{\in} \tilde{s}PO(Y)$ .

**Definition 2.18.** [1] A soft pre-open set  $(F, E)$  in a soft topological space  $(X, \tilde{\tau}, E)$  is called soft  $p_c$ -open if for each  $x_e \tilde{\in} (F, E)$ , there exists a soft closed set  $(K, E)$  such that  $x_e \tilde{\in} (K, E) \sqsubseteq (F, E)$ . The soft complement of each soft  $p_c$ -open set is called soft  $p_c$ -closed set.

The family of all soft  $p_c$ -open (resp., soft  $p_c$ -closed) sets in a soft topological space  $(X, \tilde{\tau}, E)$  is denoted by  $\tilde{s}P_cO(X, \tilde{\tau}, E)$  (resp.,  $\tilde{s}P_cC(X, \tilde{\tau}, E)$ ) or  $\tilde{s}P_cO(X)$  (resp.,  $\tilde{s}P_cC(X)$ ).

**Definition 2.19.** [2] Let  $(X, \tilde{\tau}, E)$  be a soft topological space and let  $(G, E)$  be a soft set. Then

1. The soft pre-closure of  $(G, E)$  is the soft set  $\tilde{s}pcl(G, E) = \sqcap \{(K, B) \tilde{\in} \tilde{s}PC(\tilde{X}) : (G, E) \sqsubseteq (K, B)\}$
2. The soft pre-interior of  $(G, E)$  is the soft set  $\tilde{s}pint(G, E) = \sqcup \{(H, B) \tilde{\in} \tilde{s}PO(\tilde{X}) : (H, B) \sqsubseteq (G, E)\}$ .

**Definition 2.20.** [14] Let  $(X, \tilde{\tau}, E)$  be a soft topological space and let  $(G, E)$  be a soft set. Then

1. A soft point  $x_e \tilde{\in} \tilde{X}$  is said to be a soft  $p_c$ -limit soft point of a soft set  $(F, E)$  if for every soft  $p_c$ -open set  $(G, E)$  containing  $x_e$ ,  $(G, E) \cap [(F, E) \setminus \{x_e\}] \neq \tilde{\phi}$ . The set of all soft  $p_c$ -limit soft points of  $(F, E)$  is called the soft  $p_c$ -derived set of  $(F, E)$  and is denoted by  $\tilde{s}P_cD(F, E)$ .
2. The soft  $p_c$ -closure of  $(G, E)$  is the soft set  $\tilde{s}p_ccl(G, E) = \sqcap \{(K, B) \tilde{\in} \tilde{s}P_cC(\tilde{X}) : (G, E) \sqsubseteq (K, B)\}$
3. The soft  $p_c$ -interior of  $(G, E)$  is the soft set  $\tilde{s}p_cint(G, E) = \sqcup \{(H, B) \tilde{\in} \tilde{s}P_cO(\tilde{X}) : (H, B) \sqsubseteq (G, E)\}$ .

**Lemma 2.21.** [1] If  $(F, E) \sqsubseteq \tilde{Y} \sqsubseteq \tilde{X}$  and  $\tilde{Y}$  is soft clopen. Then  $(F, E) \tilde{\in} \tilde{s}P_cO(Y)$  if and only if  $(F, E) \tilde{\in} \tilde{s}P_cO(X)$ .

**Lemma 2.22.** [1] Let  $(F, E) \sqsubseteq \tilde{Y} \sqsubseteq \tilde{X}$  and  $\tilde{Y}$  be soft clopen. If  $(F, E) \tilde{\in} \tilde{s}P_cO(X)$ , then  $(F, E) \cap \tilde{Y} \tilde{\in} \tilde{s}P_cO(Y)$ .

**Lemma 2.23.** [14] Let  $(F, E) \sqsubseteq \tilde{Y} \sqsubseteq \tilde{X}$ . If  $\tilde{Y}$  is soft clopen, then  $\tilde{s}p_ccl_Y(F, E) = \tilde{s}p_ccl_X(F, E) \cap \tilde{Y}$ .

**Definition 2.24.** [13] A soft topological space  $(X, \tilde{\tau}, E)$  is said to be:

1. Soft  $T_0$ , if for each pair of distinct soft points  $x, y \tilde{\in} X$ , there exist soft open sets  $(F, E)$  and  $(G, E)$  such that either  $x \tilde{\in} (F, E)$  and  $y \notin (F, E)$  or  $y \tilde{\in} (G, E)$  and  $x \notin (G, E)$ .

2. Soft  $T_1$ , if for each pair of distinct soft points  $x, y \in X$ , there exist two soft open sets  $(F, E)$  and  $(G, E)$  such that  $x \in (F, E)$  but  $y \notin (F, E)$  and  $y \in (G, E)$  but  $x \notin (G, E)$ .
3. Soft  $T_2$ , if for each pair of distinct soft points  $x, y \in X$ , there exist two disjoint soft open sets  $(F, E)$  and  $(G, E)$  containing  $x$  and  $y$  respectively.

In [10], S. Bayramov and C. G. Aras redefined soft  $T_i$ -spaces by replacing soft points  $x_e, y_{e'}$  instead of the ordinary points  $x, y$  in Definition 2.24.

**Proposition 2.2.** [10]

1. Every soft  $T_2$ -space  $\Rightarrow$  soft  $T_1$ -space  $\Rightarrow$  soft  $T_0$ -space.
2. A soft topological space  $(X, \tilde{\tau}, E)$  is soft  $T_1$  if and only if each soft point is soft closed.

In [24], a soft regular space is defined by using ordinary points as:

**Definition 2.25.** [24] If for every  $x \in X$  and every soft closed set  $(F, E)$  not containing  $x$ , there exist two soft open sets  $(G, E)$  and  $(H, E)$  such that  $x \in (G, E)$ ,  $(F, E) \sqsubseteq (H, E)$  and  $(G, E) \cap (H, E) = \tilde{\phi}$  then  $\tilde{X}$  is called soft regular.

In [15] a soft regular space is defined by replacing soft points  $x_e$  instead of the ordinary point  $x$  in Definition 2.25.

**Definition 2.26.** [18] A soft topological space  $(X, \tilde{\tau}, E)$  is said to be

1.  $\tilde{s}p_cT_0$ , if for each pair of distinct soft points  $x_e, y_{e'} \in SP(X)_E$ , there exist soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $x_e \in (F, E)$  and  $y_{e'} \notin (F, E)$  or  $y_{e'} \in (G, E)$  and  $x_e \notin (G, E)$ .
2.  $\tilde{s}p_cT_1$ , if for each pair of distinct soft points  $x_e, y_{e'} \in SP(X)_E$ , there exist two soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $x_e \in (F, E)$  but  $y_{e'} \notin (F, E)$  and  $y_{e'} \in (G, E)$  but  $x_e \notin (G, E)$ .
3.  $\tilde{s}p_cT_2$ , if for each pair of distinct soft points  $x_e, y_{e'} \in SP(X)_E$ , there exist two disjoint soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  containing  $x_e$  and  $y_{e'}$  respectively.

**Definition 2.27.** [18] A soft topological space  $(X, \tilde{\tau}, E)$  is said to be

1.  $\tilde{s}p_cT_0^*$ , if for each pair of distinct points  $x, y \in X$ , there exist soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $x \in (F, E)$  and  $y \notin (F, E)$  or  $y \in (G, E)$  and  $x \notin (G, E)$ .
2.  $\tilde{s}p_cT_1^*$ , if for each pair of distinct points  $x, y \in X$ , there exist two soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $x \in (F, E)$  but  $y \notin (F, E)$  and  $y \in (G, E)$  but  $x \notin (G, E)$ .

3.  $\tilde{s}p_c-T_2^*$ , if for each pair of distinct soft points  $x, y \in X$ , there exist two disjoint soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  containing  $x$  and  $y$  respectively.

**Proposition 2.3.** [18] A space  $(X, \tilde{\tau}, E)$  is  $\tilde{s}p_c-T_0$  if and only if every soft points  $x_e \neq y_{e'}$  implies  $\tilde{s}p_c cl\{x_e\} \neq \tilde{s}p_c cl\{y_{e'}\}$ .

**Proposition 2.4.** [18] A space  $(X, \tilde{\tau}, E)$  is  $\tilde{s}p_c-T_1$  if and only if every soft point of the space  $(X, \tilde{\tau}, E)$  is an soft  $p_c$ -closed set.

**Proposition 2.5.** [18] If a soft topological space  $(X, \tilde{\tau}, E)$  is  $\tilde{s}p_c-T_1$ , then it is soft  $\tilde{s}p_c-T_1^*$ .

**Definition 2.28.** [19] Let  $SP(X)_E$  and  $SP(Y)_B$  be families of soft sets. Let  $u : X \rightarrow Y$  and  $p : E \rightarrow B$  be mappings. Then a mapping  $f_{pu} : SP(X)_E \rightarrow SP(Y)_B$  is defined as:

1. Let  $(F, E)$  be a soft set in  $SP(X)_E$ . The image of  $(F, E)$  under  $f_{pu}$ , written as  $f_{pu}(F, E) = (f_{pu}(F), p(E))$ , is a soft set in  $SP(Y)_B$  such that

$$f_{pu}(F)(e') = \begin{cases} \bigcup_{e \in p^{-1}(e') \cap E} u(F(e)) & : \text{ if } p^{-1}(e') \cap E \neq \phi \\ \phi & : \text{ if } p^{-1}(e') \cap E = \phi \end{cases}$$

for all  $e' \in B$ .

2. Let  $(G, B)$  be a soft set in  $SP(Y)_B$ . Then the inverse image of  $(G, B)$  under  $f_{pu}$ , written as  $f_{pu}^{-1}(G, B) = (f_{pu}^{-1}(G), p^{-1}(B))$ , is a soft set in  $SP(X)_E$  such that

$$f_{pu}^{-1}(G)(e) = \begin{cases} u^{-1}(G(p(e))) & : \text{ if } p(e) \in B \\ \phi & : \text{ otherwise} \end{cases}$$

for all  $e \in E$ .

The soft function  $f_{pu}$  is called surjective if  $p$  and  $u$  are surjective and it is called injective if  $p$  and  $u$  are injective.

**Definition 2.29.** [25] Let  $(X, \tilde{\tau}, E)$  and  $(Y, \tilde{\mu}, B)$  be two soft topological spaces. A soft mapping  $f_{pu} : (X, \tilde{\tau}, E) \rightarrow (Y, \tilde{\mu}, B)$  is called soft continuous if  $f_{pu}^{-1}((G, B)) \tilde{\tau}$  for all  $(G, B) \tilde{\mu}$ .

### 3 Soft $p_c$ -regular spaces

In this section, we introduce some types of soft regular spaces by using soft  $p_c$ -open sets. Many characterizations of these spaces are found. Also some hereditary properties and relations between these spaces are investigated.

**Definition 3.1.** A soft space  $\tilde{X}$  is said to be  $\tilde{sp}_c$ -regular (resp.,  $\tilde{sp}_c^*$ -regular), if for each  $x_e \in \tilde{X}$  and each soft closed (resp.,  $\tilde{sp}_c$ -closed) set  $(K, E)$  such that  $x_e \notin (K, E)$ , there exist two disjoint soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $x_e \in (F, E)$  and  $(K, E) \sqsubseteq (G, E)$ .

**Remark 3.1.** In a finite soft space  $SP(X)_E$ , if  $(F, E)$  is any soft  $p_c$ -open set, then by definition it is soft pre-open and a union of soft closed sets and hence it is soft closed, so we obtain that  $(F, E)$  is both soft open and soft closed. Equivalently, any soft  $p_c$ -closed set is both open and closed.

From the above remark, we get the following result

**Proposition 3.2.** If  $SP(X)_E$  is finite, then every  $\tilde{sp}_c$ -regular space is both  $\tilde{sp}_c^*$ -regular and soft regular.

The following example shows that an  $\tilde{sp}_c$ -regular space is not necessary  $\tilde{sp}_c - T_i$  for  $i = 0, 1, 2$ .

**Example 3.2.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  and  $\tilde{X} = \{(e_1, X), (e_2, X)\}$  and let  $\tilde{\tau} = \{\tilde{X}, \tilde{\phi}, (F_1, E), (F_2, E), (F_3, E), (F_4, E), (F_5, E), (F_6, E)\}$ , where  $(F_1, E) = \{(e_1, \{x\}), (e_2, \phi)\}$ ,  $(F_2, E) = \{(e_1, \{x\}), (e_2, X)\}$ ,  $(F_3, E) = \{(e_1, \{y\}), (e_2, \phi)\}$ ,  $(F_4, E) = \{(e_1, \{y\}), (e_2, X)\}$ ,  $(F_5, E) = \{(e_1, X), (e_2, \phi)\}$ ,  $(F_6, E) = \{(e_1, \phi), (e_2, X)\}$ . Then it can be checked that  $\tilde{sp}_c O(X) = \tilde{\tau}$ . Since  $x_{e_2} \neq y_{e_2}$  and each soft open set containing one of them contains the other, so it is not  $\tilde{sp}_c - T_i$  for  $i = 0, 1, 2$ . This space is  $\tilde{sp}_c - T_i^*$  for  $i = 0, 1$  but it is not  $\tilde{sp}_c - T_2^*$ . By easy calculation it can be shown that this space is  $\tilde{sp}_c$ -regular and hence by Proposition 3.2 it is both  $\tilde{sp}_c^*$ -regular and soft regular.

**Example 3.3.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  and  $\tilde{X} = \{(e_1, X), (e_2, X)\}$  and let  $\tilde{\tau} = \{\tilde{X}, \tilde{\phi}, (F_1, E), (F_2, E), (F_3, E), (F_4, E)\}$ , where  $(F_1, E) = \{(e_1, X), (e_2, \phi)\}$ ,  $(F_2, E) = \{(e_1, \phi), (e_2, \{x, y\})\}$ ,  $(F_3, E) = \{(e_1, \{y\}), (e_2, X)\}$  and  $(F_4, E) = \{(e_1, \{y\}), (e_2, \phi)\}$ . Since  $y_{e_1} \notin (F_3, E)^c$  but there are no disjoint soft  $p_c$ -open sets containing them. Hence, this space is not  $\tilde{sp}_c$ -regular and not soft regular but it can be checked that it is  $\tilde{sp}_c^*$ -regular.

Recall that a soft space  $(X, \tilde{\tau}, E)$  is called soft-Alexandroff space [23] if any arbitrary intersection of soft open sets is soft open. Equivalently, any arbitrary union of soft closed sets is soft closed.

**Proposition 3.3.** Every soft-Alexandroff space is  $\tilde{sp}_c^*$ -regular.

*Proof.* Similar to Remark 3.1, in a soft-Alexandroff space  $(X, \tilde{\tau}, E)$ . If  $(F, E)$  is an  $\tilde{sp}_c^*$ -open set, then it is soft closed and hence  $(F, E)$  and its complement are both soft open and soft closed. Therefore, for each  $x_e \notin (F, E)$ , we have  $(F, E)$  and  $(F, E)^c$  are the required disjoint soft  $\tilde{sp}_c^*$ -open sets.  $\square$

If we take  $X = \mathbb{R}$  with the usual topology and if  $E$  consists only one parameter, then  $\mathbb{R}$  is both soft regular and  $\tilde{sp}_c^*$ -regular but it is not soft-Alexandroff.



**Theorem 3.4.** *The following statements about a space  $\tilde{X}$  are equivalent:*

1.  $\tilde{X}$  is  $\tilde{s}p_c^*$ -regular (resp.,  $\tilde{s}p_c$ -regular) space.
2. For each  $x_e \tilde{\in} \tilde{X}$  and each soft  $p_c$ -open (resp., soft open) set  $(F, E)$  containing  $x_e$ , there exist soft  $p_c$ -open set  $(G, E)$  containing  $x_e$  such that  $x_e \tilde{\in} (G, E) \sqsubseteq \tilde{s}p_c cl(G, E) \sqsubseteq (F, E)$ .
3. Each element of  $X$  has an  $\tilde{s}p_c$ -neighborhood (resp.; soft neighborhood) base consisting of soft  $p_c$ -closed sets.
4. Every soft  $p_c$ -closed (resp.; soft closed) set  $(K, E)$  is the intersection of all soft  $p_c$ -closed neighborhoods of  $(K, E)$ .
5. For every non-empty soft subset  $(F, E)$  of  $\tilde{X}$  and every soft  $p_c$ -open (resp., soft open) subset  $(G, E)$  of  $\tilde{X}$  such that  $(F, E) \cap (G, E) \neq \tilde{\phi}$ , there exist  $\tilde{s}p_c$ -open subset  $(W, E)$  of  $\tilde{X}$  such that  $(F, E) \cap (W, E) \neq \tilde{\phi}$ , and  $\tilde{s}p_c cl(W, E) \sqsubseteq (G, E)$ .
6. For every non-empty soft subset  $(F, E)$  of  $\tilde{X}$  and every soft  $p_c$ -closed (resp., soft closed) subset  $(K, E)$  of  $\tilde{X}$  such that  $(F, E) \cap (K, E) = \tilde{\phi}$ , there exist two soft  $p_c$ -open subset  $(G, E)$  and  $(W, E)$  such that  $(F, E) \cap (G, E) \neq \tilde{\phi}$ ,  $(W, E) \cap (G, E) = \tilde{\phi}$  and  $(K, E) \sqsubseteq (W, E)$ .

*Proof.* We only prove the  $\tilde{s}p_c^*$ -regular case. Since the other case can be proved similarly.

(1)  $\rightarrow$  (2). Let  $(F, E)$  be soft  $p_c$ -open set and  $x_e \tilde{\in} (F, E)$ . Then  $\tilde{X} \setminus (F, E)$  is a soft  $p_c$ -closed set such that  $x_e \tilde{\notin} \tilde{X} \setminus (F, E)$ . By  $\tilde{s}p_c^*$ -regularity of  $\tilde{X}$ , there are soft  $p_c$ -open sets  $(G, E), (H, E)$  such that  $x_e \tilde{\in} (G, E), \tilde{X} \setminus (F, E) \sqsubseteq (H, E)$  and  $(H, E) \cap (G, E) = \tilde{\phi}$ . Therefore,  $x_e \tilde{\in} (G, E) \sqsubseteq \tilde{X} \setminus (H, E) \sqsubseteq (F, E)$ . Hence  $x_e \tilde{\in} (G, E) \sqsubseteq \tilde{s}p_c cl(G, E) \sqsubseteq \tilde{s}p_c cl(\tilde{X} \setminus (H, E)) = \tilde{X} \setminus (H, E) \sqsubseteq (F, E)$ . This gives  $\tilde{s}p_c cl(G, E) \sqsubseteq \tilde{X} \setminus (H, E) \sqsubseteq (F, E)$ . Consequently,  $x_e \tilde{\in} (G, E)$  and  $\tilde{s}p_c cl(G, E) \sqsubseteq (F, E)$ .

(2)  $\rightarrow$  (3). Let  $y_{e'} \tilde{\in} \tilde{X}$ . Then for every soft  $p_c$ -open set  $(G, E)$  such that  $y_{e'} \tilde{\in} (G, E)$ ,  $\tilde{s}p_c cl(G, E) \sqsubseteq (F, E)$ . Thus for each  $y_{e'} \tilde{\in} \tilde{X}$ , the sets  $\tilde{s}p_c cl(G, E)$  form an  $\tilde{s}p_c$ -neighborhood base consisting of soft  $p_c$ -closed sets of  $\tilde{X}$ . This proves (3).

(3)  $\rightarrow$  (1). Let  $(K, E)$  be soft  $p_c$ -closed set which does not contain  $x_e$ . Then  $\tilde{X} \setminus (K, E)$  is soft  $p_c$ -open, so it is  $\tilde{s}p_c$ -neighborhood of  $x_e$ . By (3), there is soft  $p_c$ -closed set  $(L, E)$  which contains  $x_e$  and it is an  $\tilde{s}p_c$ -neighborhood of  $x_e$  with  $(L, E) \sqsubseteq \tilde{X} \setminus (K, E)$ . Consider the sets  $(L, E)$  and  $\tilde{X} \setminus (L, E)$ . Then  $x_e \tilde{\in} (L, E), (K, E) \sqsubseteq \tilde{X} \setminus (L, E) = (G, E)$  and  $(K, E) \cap (L, E) = \tilde{\phi}$ . Therefore,  $\tilde{X}$  is  $\tilde{s}p_c^*$ -regular.

(2)  $\rightarrow$  (4). Let  $(K, E)$  be soft  $p_c$ -closed and  $x_e \tilde{\notin} (K, E)$ . Then  $x_e \tilde{\in} \tilde{X} \setminus (K, E)$  and  $\tilde{X} \setminus (K, E)$  is  $\tilde{s}p_c$ -open subset of  $\tilde{X}$ . Using the hypothesis, there exists an

soft  $p_c$ -open set  $(F, E)$  such that  $x_e \tilde{\in}(F, E) \sqsubseteq \tilde{sp}_c cl(F, E) \sqsubseteq \tilde{X} \setminus (K, E)$ . Hence  $(K, E) \sqsubseteq \tilde{X} \setminus \tilde{sp}_c cl(F, E) \sqsubseteq \tilde{X} \setminus (F, E)$ . Consequently  $\tilde{X} \setminus (F, E)$  is soft  $p_c$ -closed neighborhood of  $(K, E)$  to which  $x_e$  does not belong. This proves (4).

(4)  $\rightarrow$  (5). Let  $\phi \neq (F, E) \sqsubseteq \tilde{X}$  and  $(G, E)$  be a soft  $p_c$ -open subset of  $\tilde{X}$  such that  $(F, E) \cap (G, E) \neq \phi$ . Let  $x_e \tilde{\in}(F, E) \cap (G, E)$ . Since  $x_e \notin \tilde{X} \setminus (G, E)$  and  $\tilde{X} \setminus (G, E)$  is soft  $p_c$ -closed, so there exists a soft  $p_c$ -closed neighborhood of  $\tilde{X} \setminus (G, E)$  say  $(E, E)$ , such that  $x_e \tilde{\notin}(E, E)$ . Let  $\tilde{X} \setminus (G, E) \sqsubseteq (D, E) \sqsubseteq (E, E)$  where  $(D, E)$  is soft  $p_c$ -open set. Then  $(W, E) = \tilde{X} \setminus (E, E)$  is soft  $p_c$ -open set,  $x_e \tilde{\in}(W, E)$  and  $(F, E) \cap (W, E) \neq \phi$ . Also  $\tilde{X} \setminus (D, E)$  being soft  $p_c$ -closed.  $\tilde{sp}_c cl(W, E) = \tilde{sp}_c cl(\tilde{X} \setminus (E, E)) \sqsubseteq \tilde{X} \setminus (D, E) \sqsubseteq (G, E)$ .

(5)  $\rightarrow$  (6). Let  $\phi \neq (F, E) \sqsubseteq \tilde{X}$  and  $(K, E)$  be soft  $p_c$ -closed subset of  $\tilde{X}$  such that  $(K, E) \cap (F, E) = \phi$ , then  $\tilde{X} \setminus (K, E) \cap (F, E) \neq \phi$ , and  $\tilde{X} \setminus (K, E)$  is soft  $p_c$ -open. Using (5), there exists a soft  $p_c$ -open subset  $(G, E)$  of  $\tilde{X}$  such that  $(G, E) \cap (F, E) \neq \phi$  and  $(G, E) \sqsubseteq \tilde{sp}_c cl(G, E) \sqsubseteq \tilde{X} \setminus (K, E)$ . Putting  $(W, E) = \tilde{X} \setminus \tilde{sp}_c cl(G, E)$ , then  $(K, E) \sqsubseteq (W, E) \sqsubseteq \tilde{X} \setminus (G, E)$ , and  $(W, E)$  is soft  $p_c$ -open. Hence the proof.

(6)  $\rightarrow$  (1). Let  $x_e \tilde{\notin}(K, E)$ , where  $(K, E)$  is soft  $p_c$ -closed, and let  $(F, E) = \{x_e\} \neq \phi$ , Then  $(K, E) \cap (F, E) = \tilde{\phi}$  and hence, using (6) there exist two soft  $p_c$ -open sets  $(G, E)$ , and  $(W, E)$  such that  $(W, E) \cap (G, E) = \tilde{\phi}$ ,  $(G, E) \cap (F, E) \neq \tilde{\phi}$  and  $(K, E) \sqsubseteq (W, E)$ , which implies that  $\tilde{X}$  is  $\tilde{sp}_c^*$ -regular.  $\square$

**Theorem 3.5.** *A topological space  $(X, \tau, E)$  is  $\tilde{sp}_c^*$ -regular (resp.,  $\tilde{sp}_c$ -regular) if and only if for each  $x_e \tilde{\in} \tilde{X}$  and soft  $p_c$ -closed (resp., soft closed) set  $(K, E)$  such that  $x_e \tilde{\notin}(K, E)$ , there exist soft  $p_c$ -open sets  $(G, E)$ ,  $(H, E)$  such that  $x_e \tilde{\in}(G, E)$ ,  $(K, E) \sqsubseteq (H, E)$  and  $\tilde{sp}_c cl(G, E) \cap \tilde{sp}_c cl(H, E) = \tilde{\phi}$ .*

*Proof.* We only prove the  $\tilde{sp}_c^*$ -regular case because the other case can be proved similarly.

Suppose that  $\tilde{X}$  is  $\tilde{sp}_c^*$ -regular, then for each  $x_e \tilde{\in} \tilde{X}$  and soft  $p_c$ -closed set  $(K, E)$  such that  $x_e \tilde{\notin}(K, E)$ , there exist two soft  $p_c$ -open sets  $(U, E)$  and  $(V, E)$  such that  $x_e \tilde{\in}(U, E)$ ,  $(K, E) \sqsubseteq (V, E)$  and  $(U, E) \cap (V, E) = \phi$ . Which implies that  $x_e \tilde{\in}(U, E) \sqsubseteq \tilde{X} \setminus (V, E) \sqsubseteq \tilde{X} \setminus (K, E)$ . That is  $x_e \tilde{\in}(U, E) \sqsubseteq \tilde{sp}_c cl(U, E) \sqsubseteq \tilde{X} \setminus (V, E) \sqsubseteq \tilde{X} \setminus (K, E)$ . Using Theorem 3.4(2) and the fact that  $x_e \tilde{\in}(U, E)$ , where  $(U, E)$  is soft  $p_c$ -open set there exist soft  $p_c$ -open  $(G, E)$  containing  $x_e$  such that  $x_e \tilde{\in}(G, E) \sqsubseteq \tilde{sp}_c cl(G, E) \sqsubseteq (U, E)$ . Therefore  $(K, E) \sqsubseteq (V, E) \sqsubseteq \tilde{X} \setminus \tilde{sp}_c cl(U, E) \sqsubseteq \tilde{X} \setminus (U, E) \sqsubseteq \tilde{X} \setminus \tilde{sp}_c cl(G, E)$  and  $(K, E) \sqsubseteq (V, E) \sqsubseteq \tilde{sp}_c cl(V, E) \sqsubseteq \tilde{X} \setminus (U, E)$ , Now take  $(H, E) = (V, E)$ , we get  $x_e \tilde{\in}(G, E)$ ,  $(K, E) \sqsubseteq (H, E)$  and  $\tilde{sp}_c cl(G, E) \cap \tilde{sp}_c cl(H, E) = \phi$ . This proves the necessity part. The proof of sufficiency follows directly.  $\square$

**Lemma 3.6.** *Every soft clopen subspace of an  $\tilde{sp}_c$ -regular space  $\tilde{X}$  is  $\tilde{sp}_c$ -regular.*

*Proof.* Let  $\tilde{Y}$  be a soft clopen subspace of  $\tilde{s}p_c$ -regular space  $\tilde{X}$ . Suppose that  $(H, E)$  is soft  $p_c$ -closed set in  $\tilde{Y}$  and  $y_{e'} \in \tilde{Y}$  such that  $y_{e'} \notin (H, E)$ . Then  $(H, E) = (G, E) \cap Y$ , where  $(G, E)$  is soft  $p_c$ -closed in  $\tilde{X}$ . Then  $y_{e'} \notin (G, E)$ . Since  $\tilde{X}$  is  $\tilde{s}p_c$ -regular, there exist disjoint soft  $p_c$ -open sets  $(U, E), (V, E)$  in  $\tilde{X}$  such that  $y_{e'} \in (U, E), (H, E) \subseteq (V, E)$ . Then  $(U, E) \cap Y$  and  $(V, E) \cap Y$  are disjoint soft  $p_c$ -open sets in  $\tilde{Y}$  containing  $y_{e'}$  and  $(H, E)$  respectively. This completes the proof.  $\square$

**Remark 3.7.** *If the soft space  $\tilde{X}$  is finite, then by Remark 3.1, every soft  $p_c$ -open set is both closed and open and hence we obtain that Lemma 3.4 is true for every subspace. Lemma 3.4 is true because the intersection of an soft  $p_c$ -open set in  $\tilde{X}$  with a soft clopen subspace remains an soft  $p_c$ -open set in the subspace but still we ask the following question.*

*Every soft subspace of an  $\tilde{s}p_c$ -regular space  $\tilde{X}$  is  $\tilde{s}p_c$ -regular or not ?*

**Theorem 3.8.** *Every  $\tilde{s}p_c$ -regular and  $\tilde{s}p_c - T_0$  space  $\tilde{X}$  is an  $\tilde{s}p_c - T_2$  space.*

*Proof.* Let  $x_e, y_{e'} \in \tilde{X}$  such that  $x_e \neq y_{e'}$ . Since  $\tilde{X}$  is  $\tilde{s}p_c - T_0$ , then there exists an soft  $p_c$ -open set  $(U, E)$  containing  $x_e$  but not  $y_{e'}$ . Using the hypothesis that  $\tilde{X}$  is  $\tilde{s}p_c$ -regular and since  $x_e \in (U, E)$ , so there is an soft  $p_c$ -open set  $(V, E)$ , such that  $x_e \in (V, E) \subseteq \tilde{s}p_c cl(V, E) \subseteq (U, E)$ . But  $y_{e'} \notin (U, E)$  implies that  $y_{e'} \notin \tilde{s}p_c cl(V, E)$ , then we get  $y_{e'} \in \tilde{X} \setminus \tilde{s}p_c cl(V, E)$ . Therefore, we have  $(U, E)$  and  $\tilde{X} \setminus \tilde{s}p_c cl(V, E)$  are soft  $p_c$ -open sets such that  $x_e \in (U, E), y_{e'} \in \tilde{X} \setminus \tilde{s}p_c cl(V, E)$  and  $(U, E) \cap (\tilde{X} \setminus \tilde{s}p_c cl(V, E)) = \tilde{\phi}$ . Hence the result follows.  $\square$

The proof of the following lemma is obvious.

**Lemma 3.9.** *Let  $(X, \tilde{\tau}, E)$  be an  $\tilde{s}p_c$ -regular (resp., an  $\tilde{s}p_c^*$ -regular) space and let  $(H, E)$  be a soft closed (resp., soft  $p_c$ -closed) set such that  $x_e \notin (H, E)$ , then there exists an soft  $p_c$ -open set  $(F, E)$  such that  $x_e \in (F, E)$  and  $(F, E) \cap (H, E) = \tilde{\phi}$ .*

**Proposition 3.4.** *A soft topological space is  $\tilde{s}p_c$ -regular (resp., an  $\tilde{s}p_c^*$ -regular) if and only if for each soft point  $x_e \in SP(X)_E$  and for each soft open (resp., soft  $p_c$ -open) set  $(F, E)$  containing  $x_e$ , there exists an soft  $p_c$ -open set  $(U, E)$  of  $x_e$  such that  $\tilde{s}p_c cl(U, E) \subseteq (F, E)$ .*

*Proof.* Let  $(X, \tilde{\tau}, E)$  be  $\tilde{s}p_c$ -regular space. Let  $x_e \in \tilde{X}$  and  $(F, E)$  is an soft  $p_c$ -open set containing  $x_e$ . Then,  $X \setminus (F, E)$  is an soft  $p_c$ -closed set such that  $x_e \notin X \setminus (F, E)$ . Since  $(X, \tilde{\tau}, E)$  is an  $\tilde{s}p_c$ -regular, so there exist soft  $p_c$ -open sets  $(V, E)$  and  $(U, E)$  such that  $x_e \in (U, E), X \setminus (F, E) \subseteq (V, E)$  and  $(U, E) \cap (V, E) = \tilde{\phi}$ . Thus,  $(U, E) \subseteq X \setminus (V, E)$  and hence  $\tilde{s}p_c cl(U, E) \subseteq X \setminus (V, E) \subseteq (F, E)$ .

Conversely, let  $x_e \in \tilde{X}$  and  $(H, E)$  be an soft  $p_c$ -closed set such that  $x_e \notin (H, E)$ . Then,  $X \setminus (H, E)$  is an soft  $p_c$ -open set containing  $x_e$ . So, by hypothesis there exist an soft  $p_c$ -open set  $(U, E)$  of  $x_e$  such that  $\tilde{s}p_c cl(U, E) \subseteq X \setminus (H, E)$ . Thus,  $(H, E) \subseteq X \setminus \tilde{s}p_c cl(U, E)$  and  $(U, E) \cap X \setminus \tilde{s}p_c cl(U, E) = \tilde{\phi}$ . Therefore,  $(X, \tilde{\tau}, E)$  is

$\tilde{s}p_c$ -regular.

The proof when  $(X, \tilde{\tau}, E)$  is  $\tilde{s}p_c$ -regular is analogues.  $\square$

**Definition 3.10.** A soft topological space  $(X, \tilde{\tau}, E)$  is said to be strongly  $\tilde{s}p_c^*$ -regular (resp., strongly  $\tilde{s}p_c$ -regular), if for every soft  $p_c$ -closed (resp., soft closed) set  $(H, E)$  and every point  $x \notin (H, E)$ , there exists disjoint soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $x \in (F, E)$  and  $(H, E) \sqsubseteq (G, E)$ .

**Example 3.11.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  and  $\tilde{X} = \{(e_1, X), (e_2, X)\}$  and let  $\tilde{\tau} = \{\tilde{X}, \tilde{\phi}, (F_1, E), (F_2, E)\}$ , where  $(F_1, E) = \{(e_1, \{x\}), (e_2, \{x\})\}$ ,  $(F_2, E) = \{(e_1, \{y\}), (e_2, \{y\})\}$ . Then it is not difficult to check that  $(X, \tilde{\tau}, E)$  is both strongly  $\tilde{s}p_c^*$ -regular and strongly  $\tilde{s}p_c$ -regular.

The following result is obvious.

**Proposition 3.5.** Every strongly  $\tilde{s}p_c^*$ -regular (resp., strongly  $\tilde{s}p_c$ -regular) space is  $\tilde{s}p_c^*$ -regular (resp.,  $\tilde{s}p_c$ -regular).

The converse of Proposition 3.5 is not true in general. The space in Example 3.2, is  $\tilde{s}p_c^*$ -regular and  $\tilde{s}p_c$ -regular but it is neither strongly  $\tilde{s}p_c^*$ -regular nor strongly  $\tilde{s}p_c$ -regular.

We shall prove all the results related to strongly  $\tilde{s}p_c^*$ -regular spaces and the proof of the results related to strongly  $\tilde{s}p_c$ -regular can be done in a similar way.

**Lemma 3.12.** If  $(X, \tilde{\tau}, E)$  is strongly  $\tilde{s}p_c^*$ -regular (resp., strongly  $\tilde{s}p_c$ -regular) space and  $(H, E)$  is an soft  $p_c$ -closed (resp., soft closed) set such that  $x \notin (H, E)$ , then there exists an soft  $p_c$ -open set  $(F, E)$  such that  $x \in (F, E)$  and  $(F, E) \cap (H, E) = \tilde{\phi}$ .

**Proposition 3.6.** A soft topological space  $(X, \tilde{\tau}, E)$  is strongly  $\tilde{s}p_c^*$ -regular (resp., strongly  $\tilde{s}p_c$ -regular) if and only if for each point  $x \in X$  and for each soft  $p_c$ -open (resp., soft open) set  $(F, E)$  containing  $x$ , there exists an soft  $p_c$ -open set  $(U, E)$  containing  $x$  such that  $\tilde{s}p_c cl(U, E) \sqsubseteq (F, E)$ .

*Proof.* Let  $(X, \tilde{\tau}, E)$  be a strongly  $\tilde{s}p_c^*$ -regular space. Let  $x \in X$  and  $(F, E)$  be an soft  $p_c$ -open set containing  $x$ . Then,  $X \setminus (F, E)$  is an soft  $p_c$ -closed set such that  $x \notin X \setminus (F, E)$ . Since  $(X, \tilde{\tau}, E)$  is  $\tilde{s}p_c^*$ -regular, then there exist soft  $p_c$ -open sets  $(V, E)$  and  $(U, E)$  such that  $x \in (U, E)$ ,  $X \setminus (F, E) \sqsubseteq (V, E)$  and  $(U, E) \cap (V, E) = \tilde{\phi}$ . Thus,  $(U, E) \sqsubseteq X \setminus (V, E)$  and hence  $\tilde{s}p_c cl(U, E) \sqsubseteq X \setminus (V, E) \sqsubseteq (F, E)$ .

Conversely, let  $x \in X$  and  $(H, E)$  be an soft  $p_c$ -closed set such that  $x \notin (H, E)$ . Then,  $X \setminus (H, E)$  is an soft  $p_c$ -open set containing  $x$ . So, by hypothesis there exists an soft  $p_c$ -open set  $(U, E)$  containing  $x$  such that  $\tilde{s}p_c cl(U, E) \sqsubseteq X \setminus (H, E)$ . Thus,  $(H, E) \sqsubseteq X \setminus \tilde{s}p_c cl(U, E)$  and  $(U, E) \cap X \setminus \tilde{s}p_c cl(U, E) = \tilde{\phi}$ . Therefore  $(X, \tilde{\tau}, E)$  is strongly  $\tilde{s}p_c^*$ -regular.  $\square$

**Proposition 3.7.** *Let  $(X, \tilde{\tau}, E)$  be a soft topological space and  $x \in X$ . If  $(X, \tilde{\tau}, E)$  is a strongly  $\tilde{s}p_c^*$ -regular (resp., strongly  $\tilde{s}p_c$ -regular) space, then the following statements are true:*

1.  $x \notin \tilde{\mathcal{H}}(H, E)$  if and only if  $(x, E) \cap (H, E) = \tilde{\phi}$  for every soft  $p_c$ -closed (resp., soft closed) set  $(H, E)$ .
2.  $x \notin \tilde{\mathcal{F}}(F, E)$  if and only if  $(x, E) \cap (F, E) = \tilde{\phi}$  for every soft  $p_c$ -open (resp., soft open) set  $(F, E)$ .

*Proof.* (1) Let  $x \notin \tilde{\mathcal{H}}(H, E)$ , then by Lemma 3.12, there exists an  $\tilde{s}p_c$ -open set  $(F, E)$  such that  $x \in \tilde{\mathcal{H}}(F, E)$  and  $(F, E) \cap (H, E) = \tilde{\phi}$ . Since  $(x, E) \sqsubseteq (F, E)$ , we have  $(x, E) \cap (H, E) = \tilde{\phi}$ .

Conversely, straightforward.  $\square$

(2) Let  $x \notin \tilde{\mathcal{F}}(F, E)$ . Then we have two cases :

- (i)  $x \notin F(\alpha)$  for all  $e \in E$ , it is obvious that  $(x, E) \cap (F, E) = \tilde{\phi}$ .
- (ii)  $x \notin F(\alpha)$  and  $x \in F(\beta)$  for some  $\alpha, \beta \in E$ , then we have  $x \in X \setminus F(\alpha)$  and  $x \notin \tilde{X} \setminus F(\beta)$  for some  $\alpha, \beta \in E$  and so  $\tilde{X} \setminus (F, E)$  is an soft  $p_c$ -closed set such that  $x \notin \tilde{X} \setminus (F, E)$ , by (1),  $(x, E) \cap \tilde{X} \setminus (F, E) = \tilde{\phi}$ . So,  $(x, E) \sqsubseteq (F, E)$  but this contradicts that  $x \notin F(\alpha)$  for some  $\alpha \in E$ . Consequently, we have  $(x, E) \cap (F, E) = \tilde{\phi}$ .

The converse part is obvious.  $\square$

**Proposition 3.8.** *Let  $(X, \tilde{\tau}, E)$  be a soft topological space and  $x \in X$ . Then the following statements are equivalent:*

1.  $(X, \tilde{\tau}, E)$  is a strongly  $\tilde{s}p_c^*$ -regular (resp., strongly  $\tilde{s}p_c$ -regular) space,
2. For each soft  $p_c$ -closed (resp., soft closed) set  $(H, E)$  such that  $(x, E) \cap (H, E) = \tilde{\phi}$ , there exist soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $(x, E) \sqsubseteq (F, E)$ ,  $(H, E) \sqsubseteq (G, E)$  and  $(F, E) \cap (G, E) = \tilde{\phi}$ .

*Proof.* Follows from Proposition 3.7(1) and Lemma 3.12.  $\square$

**Proposition 3.9.** *Let  $(X, \tilde{\tau}, E)$  be a soft topological space and  $x \in X$ . If  $(X, \tilde{\tau}, E)$  is a strongly  $\tilde{s}p_c^*$ -regular (resp., strongly  $\tilde{s}p_c$ -regular), then the following statements are true:*

1. For an soft  $p_c$ -open (resp., soft open) set  $(F, E)$ ,  $x \in \tilde{\mathcal{H}}(F, E)$  if and only if  $x \in F(\alpha)$  for some  $\alpha \in E$ .
2. For an soft  $p_c$ -open (resp., soft open) set  $(F, E)$ ,  $(F, E) = \sqcup \{(x, E) : x \in F(\alpha)\}$  for some  $\alpha \in E$ .

*Proof.* (1). Suppose that  $x \in F(\alpha)$  and  $x \notin \tilde{\mathcal{H}}(F, E)$  for some  $\alpha \in E$ . Then by Proposition 3.8(2),  $(x, E) \cap (F, E) = \tilde{\phi}$ . By our assumption, this is a contradiction and so  $x \in \tilde{\mathcal{H}}(F, E)$ . The Converse is obvious.

(2). Follows from part (1) and Remark 2.8.  $\square$

**Proposition 3.10.** *Let  $(X, \tilde{\tau}, E)$  be a soft topological space and  $x \in X$ . If  $(X, \tilde{\tau}, E)$  is strongly  $\tilde{s}p_c^*$ -regular, then the following statements are equivalent:*

1.  $(X, \tilde{\tau}, E)$  is a  $\tilde{s}p_c-T_1^*$  space,
2. For  $x, y \in X$  with  $x \neq y$ , there exist soft  $p_c$ -open sets  $(F, E)$  and  $(G, E)$  such that  $(x, E) \sqsubseteq (F, E)$  and  $(y, E) \cap (F, E) = \tilde{\phi}$ ,  $(y, E) \sqsubseteq (G, E)$  and  $(x, E) \cap (G, E) = \tilde{\phi}$ .

*Proof.* It is clear that  $x \tilde{\in} (F, E)$  if and only if  $(x, E) \sqsubseteq (F, E)$ , and by Proposition 3.9(2),  $x \tilde{\notin} (F, E)$  if and only if  $(x, E) \cap (F, E) = \tilde{\phi}$ . Hence, statements (1) and (2) are equivalent.  $\square$

## 4 Soft $p_c$ -normal spaces

In this section, we define  $\tilde{s}p_c$ -normal spaces and derive many of its properties. The relationship to other soft spaces and its image under  $\tilde{s}p_c$ -continuous functions are discussed.

**Definition 4.1.** *A soft space  $\tilde{X}$  is said to be  $\tilde{s}p_c$ -normal (resp.,  $\tilde{s}p_c^*$ -normal) space, if for any disjoint soft closed (resp.,  $\tilde{s}p_c^*$ -closed) sets  $(K, E)$  and  $(L, E)$  of  $\tilde{X}$ , there exist soft  $p_c$ -open sets  $(U, E)$ ,  $(V, E)$  such that  $(K, E) \sqsubseteq (U, E)$ ,  $(L, E) \sqsubseteq (V, E)$  and  $(V, E) \cap (U, E) = \tilde{\phi}$ .*

**Example 4.2.** Let  $X = \{x_1, x_2, x_3\}$ ,  $E = \{e_1, e_2\}$  and let  $\tilde{\tau} = \{\tilde{X}, \tilde{\phi}, (F_1, E), (F_2, E), (F_3, E), (F_4, E)\}$ , where  $(F_1, E) = \{(e_1, \{x_1, x_2\}), (e_2, \{x_3\})\}$ ,  $(F_2, E) = \{(e_1, \{x_3\}), (e_2, \{x_1, x_2\})\}$ ,  $(F_3, E) = \{(e_1, \{x_1\}), (e_2, \phi)\}$ ,  $(F_4, E) = \{(e_1, \{x_1, x_3\}), (e_2, \{x_1, x_2\})\}$ . Then this space is both  $\tilde{s}p_c$ -normal and  $\tilde{s}p_c^*$ -normal but it is not  $\tilde{s}p_c$ -regular.

**Theorem 4.3.** *A space  $\tilde{X}$  is an  $\tilde{s}p_c^*$ -normal space, if for each pair of soft  $p_c$ -open sets  $(U, E)$  and  $(V, E)$  in  $\tilde{X}$  such that  $\tilde{X} = (U, E) \sqcup (V, E)$ , there are soft  $p_c$ -closed sets  $(G, E)$  and  $(H, E)$  which are contained in  $(U, E)$  and  $(V, E)$  respectively and  $\tilde{X} = (G, E) \sqcup (H, E)$ .*

*Proof.* Straightforward.  $\square$

**Theorem 4.4.** *If  $\tilde{X}$  is any soft space, then the following statements are equivalent:*

1.  $\tilde{X}$  is  $\tilde{s}p_c^*$ -normal,
2. For each  $\tilde{s}p_c$ -closed set  $(F_1, E)$  in  $\tilde{X}$  and soft  $p_c$ -open set  $(G, E)$  contains  $(F_1, E)$ , there is an soft  $p_c$ -open set  $(U, E)$  such that  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_c \text{cl}(U, E) \sqsubseteq (G, E)$ ,
3. For each  $\tilde{s}p_c$ -closed set  $(F_1, E)$  in  $\tilde{X}$  and soft  $p_c$ -open set  $(G, E)$  containing  $(F_1, E)$ , there are soft  $p_c$ -open sets  $(U_n, E)$  for  $n \in \mathbb{N}$ , such that  $(F_1, E) \sqsubseteq \bigsqcup_{n \in \mathbb{N}} (U_n, E)$  and  $\tilde{s}p_c \text{cl}(U_n, E) \sqsubseteq (G, E)$ , for each  $n \in \mathbb{N}$ .

*Proof.* (1)  $\rightarrow$  (2). Since  $(G, E)$  is soft  $p_c$ -open set containing  $(F_1, E)$ , then  $\tilde{X} \setminus (G, E)$  and  $(F_1, E)$  are disjoint soft  $p_c$ -closed sets in  $\tilde{X}$ . Since  $\tilde{X}$  is  $\tilde{s}p_c^*$ -normal, so there exist soft  $p_c$ -open sets  $(U, E)$  and  $(V, E)$  such that  $(F_1, E) \sqsubseteq (U, E)$ ,  $\tilde{X} \setminus (G, E) \sqsubseteq (V, E)$  and  $(V, E) \cap (U, E) = \tilde{\phi}$ . Hence,  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_ccl(U, E) \sqsubseteq \tilde{s}p_ccl(\tilde{X} \setminus (V, E)) = \tilde{X} \setminus (V, E) \sqsubseteq (G, E)$ , or  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_ccl(U, E) \sqsubseteq (G, E)$ .

(2)  $\rightarrow$  (3). Let  $(F_1, E)$  be an soft  $p_c$ -closed set and  $(G, E)$  be an soft  $p_c$ -open set in an  $\tilde{s}p_c^*$ -normal space  $\tilde{X}$  such that  $(F_1, E) \sqsubseteq (G, E)$ . So by hypothesis, there is an soft  $p_c$ -open set  $(U, E)$  such that  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_ccl(U, E) \sqsubseteq (G, E)$ . If we put  $(U_n, E) = (U, E)$  for all  $n \in N$ , the proof follows.

(3)  $\rightarrow$  (1). Let  $(F_1, E)$  and  $(F_2, E)$  be a pair of disjoint soft  $p_c$ -closed set in the space  $\tilde{X}$ , then  $\tilde{X} \setminus (F_2, E)$  is an soft  $p_c$ -open set in  $\tilde{X}$  containing  $(F_1, E)$ . So by hypothesis, there are soft  $p_c$ -open sets  $(U_n, E)$  for  $n \in N$  such that

$$(F_1, E) \sqsubseteq \bigsqcup_{n \in N} (U_n, E)$$

and  $\tilde{s}p_ccl(U_n, E) \sqsubseteq \tilde{X} \setminus (F_2, E)$  for each  $n \in N$ . Since  $\tilde{X} \setminus (F_1, E)$  is an soft  $p_c$ -open subset of  $\tilde{X}$  containing the soft  $p_c$ -closed set  $(F_2, E)$ , then by applying the condition of the theorem again, we get soft  $p_c$ -open sets  $(V_n, E)$  for  $n \in N$ , such that

$$(F_2, E) \sqsubseteq \bigsqcup_{n \in N} (V_n, E)$$

and  $\tilde{s}p_ccl(V_n, E) \sqsubseteq \tilde{X} \setminus (F_1, E)$  for each  $n \in N$ . Thus  $\tilde{s}p_ccl(U_n, E) \cap (F_2, E) = \tilde{\phi}$  and  $\tilde{s}p_ccl(V_n, E) \cap (F_1, E) = \tilde{\phi}$  for each  $n \in N$ . Setting

$$(G_n, E) = (U_n, E) \setminus \bigsqcup_{n \in N} \tilde{s}p_ccl(V_n, E)$$

and

$$(H_n, E) = (V_n, E) \setminus \bigsqcup_{n \in N} \tilde{s}p_ccl(U_n, E).$$

Then

$$(U, E) = \bigsqcup_{n \in N} (G_n, E)$$

and

$$(V, E) = \bigsqcup_{n \in N} (H_n, E)$$

are disjoint soft  $p_c$ -open sets in  $\tilde{X}$  containing  $(F_1, E)$  and  $(F_2, E)$  respectively. Hence  $\tilde{X}$  is  $\tilde{s}p_c^*$ -normal.  $\square$

**Theorem 4.5.** *A soft topological space  $\tilde{X}$  is  $\tilde{s}p_c$ -normal if and only if for each soft closed set  $(F_1, E)$  in  $\tilde{X}$  and soft open set  $(G, E)$  contains  $(F_1, E)$ , there is an soft  $p_c$ -open set  $(U, E)$  such that  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_ccl(U, E) \sqsubseteq (G, E)$ .*

*Proof.* Let  $(F_1, E)$  be any soft close subset in an  $\tilde{s}p_c$ -normal space  $\tilde{X}$  and  $(G, E)$  be any soft open subset of  $\tilde{X}$  containing  $(F_1, E)$ . Then  $\tilde{X} \setminus (G, E)$  is closed and  $\tilde{X} \setminus (G, E) \cap (F_1, E) = \tilde{\phi}$ . Hence by hypothesis, there exist two disjoint soft  $p_c$ -open sets  $(U, E)$  and  $(V, E)$  such that  $(F_1, E) \sqsubseteq (U, E)$ ,  $\tilde{X} \setminus (G, E) \sqsubseteq (V, E)$  and  $(V, E) \cap (U, E) = \tilde{\phi}$ . Since  $(V, E) \cap (U, E) = \tilde{\phi}$ , then  $(U, E) \sqsubseteq \tilde{X} \setminus (V, E)$ . But  $\tilde{X} \setminus (G, E) \sqsubseteq (V, E)$ , then  $\tilde{X} \setminus (V, E) \sqsubseteq (G, E)$  and so  $(U, E) \sqsubseteq (G, E)$ . And since  $(U, E)$  and  $(V, E)$  are soft  $p_c$ -open sets, then  $\tilde{X} \setminus (V, E)$  and  $\tilde{X} \setminus (U, E)$  are soft  $p_c$ -closed sets and so  $\tilde{s}p_ccl(\tilde{X} \setminus (V, E)) = \tilde{X} \setminus (V, E)$  and  $\tilde{s}p_ccl(\tilde{X} \setminus (U, E)) = \tilde{X} \setminus (U, E)$  and then  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_ccl(U, E) \sqsubseteq \tilde{s}p_ccl(\tilde{X} \setminus (V, E)) = \tilde{X} \setminus (V, E) \sqsubseteq (G, E)$ . Thus  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_ccl(U, E) \sqsubseteq (G, E)$ .

Conversely, let the condition be satisfied and let  $(F_1, E)$ ,  $(F_2, E)$  be two disjoint soft closed subsets of  $\tilde{X}$ . Then  $(F_1, E) \sqsubseteq \tilde{X} \setminus (F_2, E)$  and since  $(F_2, E)$  is soft closed then  $\tilde{X} \setminus (F_2, E)$  is a soft open subset containing  $(F_1, E)$ . So by hypothesis, there exist soft  $p_c$ -open sets  $(U, E)$  such that  $(F_1, E) \sqsubseteq (U, E) \sqsubseteq \tilde{s}p_ccl(U, E) \sqsubseteq \tilde{X} \setminus (F_2, E)$ . Putting  $(V, E) = \tilde{X} \setminus \tilde{s}p_ccl(U, E)$ , then there exist two disjoint soft  $p_c$ -open sets  $(U, E)$  and  $(V, E)$  such that  $(F_1, E) \sqsubseteq (U, E)$  and  $(F_2, E) \sqsubseteq (V, E)$ . Therefore,  $\tilde{X}$  is  $\tilde{s}p_c$ -normal.  $\square$

**Theorem 4.6.** *Every soft  $T_1$ ,  $\tilde{s}p_c$ -normal space  $\tilde{X}$  is  $\tilde{s}p_c$ -regular.*

*Proof.* Let  $(F_1, E)$  be any soft closed subset in an  $\tilde{s}p_c$ -normal space  $\tilde{X}$  and  $x_e \in \tilde{X}$  such that  $x_e \notin (F_1, E)$ . Since  $\tilde{X}$  is soft  $T_1$  space, then  $\{x_e\}$  is soft closed subset in  $\tilde{X}$  with  $\{x_e\} \cap (F_1, E) = \tilde{\phi}$ . By  $\tilde{s}p_c$ -normality of  $\tilde{X}$ , there exist two disjoint soft  $p_c$ -open sets  $(U, E)$  and  $(V, E)$  of  $\tilde{X}$  such that  $\{x_e\} \sqsubseteq (U, E)$ , so  $x_e \in (U, E)$ ,  $(F_1, E) \sqsubseteq (V, E)$  and  $(U, E) \cap (V, E) = \tilde{\phi}$ . Thus  $\tilde{X}$  is an  $\tilde{s}p_c$ -regular space.  $\square$

**Theorem 4.7.** *If  $\tilde{Y}$  is a soft clopen subspace of an  $\tilde{s}p_c$ -normal (resp.,  $\tilde{s}p_c^*$ -normal) space  $\tilde{X}$ , then  $\tilde{Y}$  is  $\tilde{s}p_c$ -normal (resp.,  $\tilde{s}p_c^*$ -normal).*

*Proof.* Let  $\tilde{X}$  be an  $\tilde{s}p_c^*$ -normal space and  $\tilde{Y}$  be a soft clopen subspace of  $\tilde{X}$ . Let  $(K_1, E)$  and  $(K_2, E)$  be two disjoint soft  $p_c$ -closed subsets of  $\tilde{Y}$ , then By Lemma 2.23,  $(K_1, E)$  and  $(K_2, E)$  are two disjoint soft  $p_c$ -closed subsets of  $\tilde{X}$ . By  $\tilde{s}p_c^*$ -normality of  $\tilde{X}$ , there exist two soft  $p_c$ -open sets  $(F_1, E)$  and  $(F_2, E)$  such that  $(K_1, E) \sqsubseteq (F_1, E)$ ,  $(K_2, E) \sqsubseteq (F_2, E)$  and  $(F_1, E) \cap (F_2, E) = \tilde{\phi}$ , then  $(K_1, E) \sqsubseteq (F_1, E) \cap \tilde{Y}$  and  $(K_2, E) \sqsubseteq (F_2, E) \cap \tilde{Y}$ . It follows from,  $(F_1, E) \cap (F_2, E) = \tilde{\phi}$ , that  $((F_1, E) \cap \tilde{Y}) \cap ((F_2, E) \cap \tilde{Y}) = \tilde{\phi}$  and By Lemma 2.21, we have  $((F_1, E) \cap \tilde{Y})$  and  $((F_2, E) \cap \tilde{Y})$  are soft  $p_c$ -open subsets of  $\tilde{Y}$ . Hence,  $\tilde{Y}$  is  $\tilde{s}p_c^*$ -normal.  $\square$



The following example shows that Theorem 4.7, is not true when  $\tilde{Y}$  is soft open or soft closed.

**Example 4.8.** Let  $X = \{x, y, z\}$ ,  $E = \{e_1, e_2\}$  and  $\tilde{X} = \{(e_1, X), (e_2, X)\}$ , let  $\tilde{\tau} = \{\tilde{X}, \tilde{\phi}, (F_1, E), (F_2, E), (F_3, E), (F_4, E)\}$ , where  $(F_1, E) = \{(e_1, \{x\}), (e_2, X)\}$ ,  $(F_2, E) = \{(e_1, \{y\}), (e_2, \{y\})\}$ ,  $(F_3, E) = \{(e_1, \phi), (e_2, \{y\})\}$ ,  $(F_4, E) = \{(e_1, \{x, y\}), (e_2, X)\}$ . Then  $(X, \tilde{\tau}, E)$  is both  $\tilde{s}p_c^*$ -normal and  $\tilde{s}p_c$ -normal space and the soft open set  $(F_4, E)$  is not  $\tilde{s}p_c$ -normal. Also  $(X, \tilde{\tau}^c, E)$  is both  $\tilde{s}p_c^*$ -normal and  $\tilde{s}p_c$ -normal space and the soft closed set  $(F_4, E)$  is not  $\tilde{s}p_c$ -normal.

**Theorem 4.9.** *Every  $\tilde{s}p_c^*$ -normal  $\tilde{s}p_c - T_2$  space  $\tilde{X}$  is  $\tilde{s}p_c^*$ -regular.*

*Proof.* Suppose that  $(F_1, E)$  is an soft  $p_c$ -closed set and  $x_e \notin (F_1, E)$  for each  $x_e \in \tilde{X}$ . Since  $\tilde{X}$  is an  $\tilde{s}p_c - T_2$  space. Therefore by Theorem 2.4, each  $\{x_e\}$  is soft  $p_c$ -closed in  $\tilde{X}$ . Since  $\tilde{X}$  is  $\tilde{s}p_c^*$ -normal, so there exist soft  $p_c$ -open sets  $(U, E), (V, E)$  such that  $\{x_e\} \subseteq (U, E)$ ,  $(F_1, E) \subseteq (V, E)$  and  $(U, E) \cap (V, E) = \tilde{\phi}$ , this implies that  $\tilde{X}$  is  $\tilde{s}p_c^*$ -regular.  $\square$

**Definition 4.10.** *A soft mapping  $f_{pu} : (X, \tilde{\tau}, E) \rightarrow (Y, \tilde{\mu}, B)$  is called an soft  $p_c$ -open mapping if and only if the image of every soft  $p_c$ -open set in  $\tilde{X}$  is an soft  $p_c$ -open in  $Y$ .*

**Proposition 4.1.** *Let  $(X, \tilde{\tau}, E)$  and  $(Y, \tilde{\mu}, B)$  be soft topological spaces and  $f_{pu} : SP(X)_E \rightarrow SP(Y)_B$  be a soft bijective and soft  $p_c$ -open mapping. If  $(X, \tilde{\tau}, E)$  is  $\tilde{s}p_c - T_i$ , then  $(Y, \tilde{\mu}, B)$  is  $\tilde{s}p_c - T_i$  spaces ( $i = 0, 1, 2$ ).*

*Proof.* We prove only the case for  $\tilde{s}p_c - T_0$  space and the proof of the other are similar. Let  $y_{\beta 1}, y_{\beta 2} \in SP(Y)_B$  be two distinct soft points. Since  $f_{pu}$  is bijective, there exist distinct soft points  $x_{e 1}, x_{e 2} \in \tilde{X}$  such that  $f_{pu}(x_{e 1}) = y_{\beta 1}$ ,  $f_{pu}(x_{e 2}) = y_{\beta 2}$ . Since  $(X, \tilde{\tau}, E)$  is an  $\tilde{s}p_c - T_0$  space, there exist soft  $p_c$ -open sets  $(F, E), (G, E)$  such that  $x_{e 1} \in (F, E)$  and  $x_{e 2} \notin (F, E)$  or  $x_{e 2} \in (G, E)$  and  $x_{e 1} \notin (G, E)$ . As  $f_{pu}$  is an soft  $p_c$ -open mapping, then  $f_{pu}(F, E), f_{pu}(G, E)$  are soft  $p_c$ -open sets such that  $y_{\beta 1} \in f_{pu}(F, E)$  and  $y_{\beta 2} \notin f_{pu}(F, E)$  or  $y_{\beta 2} \in f_{pu}(G, E)$  and  $y_{\beta 1} \notin f_{pu}(G, E)$ . This implies that,  $(Y, \tilde{\mu}, B)$  is  $\tilde{s}p_c - T_0$ .  $\square$

**Definition 4.11.** *A function  $f_{pu} : (X, \tilde{\tau}, E) \rightarrow (Y, \tilde{\mu}, B)$  is injective soft point  $\tilde{s}p_c$ -closure if and only if for every  $x_e, y_{e'} \in \tilde{X}$  such that  $\tilde{s}p_c cl(\{x_e\}) \neq \tilde{s}p_c cl(\{y_{e'}\})$ , then  $\tilde{s}p_c cl(\{f(x_e)\}) \neq \tilde{s}p_c cl(\{f(y_{e'})\})$ .*

It is clear that the identity function from any soft topological space onto itself is a function which satisfies Definition 4.11.

**Theorem 4.12.** *If a function  $f_{pu} : (X, \tilde{\tau}, E) \rightarrow (Y, \tilde{\mu}, B)$  is injective soft point  $\tilde{s}p_c$ -closure and  $\tilde{X}$  is an  $\tilde{s}p_c - T_0$  space, then  $f_{pu}$  is soft injective.*

*Proof.* Let  $x_e, y_{e'} \in \tilde{X}$  with  $x_e \neq y_{e'}$ . Since  $\tilde{X}$  is  $\tilde{s}p_c$ - $T_0$ , therefore by Proposition 2.3,  $\tilde{s}p_c cl(\{x_e\}) \neq \tilde{s}p_c cl(\{y_{e'}\})$ . But  $f_{pu}$  is  $(1 - 1)$  soft point  $\tilde{s}p_c$ -closure, implies that  $\tilde{s}p_c cl(\{f(x_e)\}) \neq \tilde{s}p_c cl(\{f(y_{e'})\})$ . Hence  $f_{pu}(x_e) \neq f_{pu}(y_{e'})$ . Thus  $f_{pu}$  is soft injective.  $\square$

## 5 Conclusion

Many topological notions are extended to the soft topology after introducing the concept of soft topological spaces. Several classes of soft sets are defined and applied to present many notions in soft topology. In this paper, we employ the notion of soft  $p_c$ -open set to introduce some types of soft regular and soft normal spaces and give many properties of these spaces. Also we discuss relations between these spaces, hereditary properties and their images under soft  $p_c$ -continuous mappings.

## References

- [1] N. K. Ahmed and Q. H. Hamko, soft  $p_c$ -open sets and  $\tilde{s}p_c$ -continuity in soft topological spaces, ZANCO Journal of Pure and Applied Sciences, 30(6) (2017) 72–84. Crossref, Google Scholar.
- [2] M. Akdag, and A. Ozkan, On Soft Preopen Sets and Soft Pre Separation Axioms, Gazi University Journal of Science, 27(4)(2014) 1077–1083. Google Scholar
- [3] T. M. Al-shami, Corrigendum to "On soft topological space via semi-open and semi-closed soft sets, Kyungpook Mathematical Journal, 54 (2014) 221-236", Kyungpook Mathematical Journal, (2018) 58 (3) (2018) 583-588.
- [4] T. M. Al-shami, M. E. El-Shafei and M. Abo-Elhamayel, On soft topological ordered spaces, Journal of King Saud University-Science, (2018) Crossref.
- [5] T. M. Al-shami and L. D. R. Kocinac, The equivalence between the enriched and extended soft topologies, Appl. Comput. Math., 18 (2) (2019) 149–162.
- [6] T. M. Al-shami, Comments on "Soft mappings spaces", The Scientific World Journal, Volume 2019, Article ID 6903809, 2 pages.
- [7] T. M. Al-shami and M. E. El-Shafei, Partial belong relation on soft separation axioms and decision-making problem, two birds with one stone, Soft Comput., (2019), Crossref.

- [8] T. M. Al-shami and M. E. El-Shafei, Two new types of separation axioms on supra soft separation spaces, *Demonstratio Mathematica*, 52 (1) (2019) 147-165.
- [9] T. M. Al-shami and M. E. El-Shafei, On supra soft topological ordered spaces, *Arab Journal of Basic and Applied Sciences*, 26 (1) (2019) 433-445
- [10] S. Bayramov and C. G. Aras, A new approach to separability and compactness in soft topological spaces, *(TWMS) Journal of Pure and Applied Mathematics* 9(1) (2018) 82–93. Google Scholar.
- [11] M. E. El-Shafei, M. Abo-Elhamayel and T. M. Al-Shami, Partial soft separation axioms and soft compact spaces, *Filomat* 32:13(2018) 4755–4771. Crossref, Google Scholar.
- [12] D. N. Georgiou, A. C. Megaritis and V. I. Petropoulos, On soft topological spaces, *Applied Mathematics and Information Sciences* 7(2) (2013) 1889-1901. Google Scholar.
- [13] O. Gocur and A. Kopuzlu, On soft separation axioms, *Annals of Fuzzy Mathematics and Informatics* 9(5)(2015) 817–822. Google Scholar.
- [14] Q. H. Hamko and N. K. Ahmed, Characterizations of soft  $p_c$ -open Sets and  $\tilde{s}p_c$ -almost continuous mapping in soft topological spaces, accepted for publication.
- [15] S. Hussain and B Ahmad, Some properties of soft topological spaces, *Computers and Mathematics with Applications* 62(2011) 4058—4067. Google Scholar.
- [16] S. Hussain and B. Ahmad , Soft separation axioms in soft topological spaces, *Hacettepe Journal of Mathematics and Statistics* 44(3) (2015) 559–568. Google Scholar.
- [17] G. Ilango and M. Ravindran, On soft pre-open sets in soft topological spaces, *International Journal of Mathematics Research* 5(4) (2013) 399–409. Google Scholar.
- [18] Alias B. Khalaf, Q. H. Hamko and N. K. Ahmed, On soft  $p_c$ -separation axioms, submitted for publication.
- [19] A. Kharal and B. Ahmad, Mappings on soft classes, *New Mathematics and Natural Computation* 7(3) (2011) 471–481. Crossref, Google Scholar.

- [20] P. K. Maji, R. Biswas and R. Roy, Soft set theory, *Computers and Mathematics with Applications* 45(2003) 555–562. Google Scholar.
- [21] D. Molodtsov, Soft set theory-first results, *Computers and Mathematics with Applications* 37 (1999) 19–31. Google Scholar.
- [22] M. Ravindran and G. Ilango, A note on soft pre-pen sets, *International Journal of Pure and Applied Mathematics* 106(5) (2016) 63–78. Crossref
- [23] A. Selvi I. Arockiarani, Soft Alexandroff spaces in soft ideal topological spaces, *Indian Journal of Applied Research* 6(3) (2016) 233–243.
- [24] M. Shabir and M. Naz, On soft topological spaces, *Computers and Mathematics with Applications* 61 (2011) 1786–1799. Crossref.
- [25] I. Zorlutuna and H. Cakır, On Continuity of Soft Mappings, *Applied Mathematics and Information Sciences* 9(1) (2015) 403–409. Google Scholar.