

**Salahaddin University-Erbil**  
**College of Engineering**  
**Department of Water Resources Engineering**  
**2022/2023**



# **Mathematics II**

## **Chapter seven**

# **Products and powers of Trigonometric Functions**

Shawnm Mudhafar Saleh

[shawnm.saleh@su.edu.krd](mailto:shawnm.saleh@su.edu.krd)

Since

- $\int \sin^n(ax) \cos ax \, dx = \frac{\sin^{n+1} ax}{a(n+1)} + c$  for  $n \neq -1$

- If  $n = -1$

- $\int \frac{\cos ax}{\sin ax} \, dx = \frac{1}{a} \ln|\sin ax| + c$

- And

- $\int \cos^n(ax) \sin ax \, dx = -\frac{\cos^{n+1} ax}{a(n+1)} + c$  for  $n \neq -1$

- $\int \tan ax \, dx = \int \frac{\sin ax}{\cos ax} = -\frac{1}{a} \ln|\cos ax| + c$

## Power condition

- $\int \sin^3 x \, dx$
- $\int \cos^{2n+1} x \, dx$
- $\int \sin^{2n+1} x \, dx$
- $\int \tan^4 x \, dx$
- $\int \tan^n x \, dx$
- $\int \sec^{2n} x \, dx$
- $\int \sec^6 x \, dx$