#### **Poultry production**

Grade 3

the science that seeks to provide learners with knowledge on essential building and equipment, incubation of eggs, hatchery management, principles for successful production, breeds and breeding, brooding of chicks and management techniques, how to rear chicks, table egg and meat production, processing and marketing, and products, health management practices, diseases and parasites, and economic implication of these diseases.

Poultry production began to be more specialized in the late 1800s and today is a highly specialized industry. The poultry industry is also actively promoting and marketing products both to meet consumer needs and to address concerns related to health and the safety of food products.

**Poultry** are domesticated birds kept by humans including (chickens; turkeys; geese; ducks; gamebirds; bantams... etc.) for their eggs, meat and feather.

# Advantages of Poultry Raising

Raising poultry has a number of advantages, among which are:

- 1. High feed efficiency.
- 2. Fast return on investment.
- 3. Spreading income throughout the year.
- 4. High return compared to feed costs.
- 5. Low land requirements.
- 6. Adaptability to both small part-time enterprises and large commercial enterprises.
- 7. The operation can be highly mechanized, with high output per hour of labor.
- 8. Getting benefits of their products.

### **Poultry products:**

1. Poultry meat: either marketed as a live bird or they are converted to different poultry meat products.

Grade 3

- 2. Eggs: either consumed as whole egg or processed in egg processing industrial.
- 3. Feather for industry and ornamental.
- 4. Poultry waste (including the wastes of poultry houses (manure) and slaughter houses) is an excellent source of organic materials which can be utilized as organic fertilizer in growing crops and concentrated feed in livestock production.

# **Classification of Poultry**

#### A. The scientific classification of chicken

- Kingdom: Animalia
- Phylum: Chordata
- Class:
- Sub-Class: Neornithes
- Order: Galliformes
- Family: Phasianidae
- Phasianinae • Subfamily:
- Genus: Gallus
- Species: G. gallus
- Subspecies: G. g. domesticus
- Scientific Name: Gallus gallus domesticus

# **B.** Classification According to Origin

According to origin the chickens are of four types:

- 1. **Asiatic**: Brahma, Longson, Cochin, Asil etc.
- 2. **English**: Austrolorp, Cornish, Dorking, Orpington etc.
- 3. Mediterranian: Leghorn, Minorca, Ancona, Fayoumi etc.
- 4. **American**: Road island red, New Hampshire, Plymouth Rock etc.

**Table: Characteristics of Some Typical Breeds of Chickens** 

Minorca (Black) Andalusian (Blue)  White White Single Brown White Dark slate Dark slate Black in size and are used mainly for egg production of crossbreeding.  American Plymouth Rock (Barred)  Brown Yellow Single Reddish bay Red Yellow Barred (sex-linked)  Plymouth Rock (White)  Brown Yellow Single Reddish bay Red Yellow White Primary use—Broiler.  New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English  Cornish (White)  Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; bottom feet white Used in production of crossbreeds.  Asiatic  Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in crossbreeding.	Class and Breed	Eggs	Skin	Comb	Eyes	Earlobes	Shanks	Plumage	Comments
Minorca (Black) White Andalusian (Blue) White White Single Brown White Andalusian (Blue) White White Single Reddish bay White Dark slate Black in size and are used mainly for egg production of corsostereding.  American Plymouth Rock (Barred) Brown Yellow Single Reddish bay Red Yellow Barred (sex-linked) Dual purpose; used in crosser sexing chicks a hatching.  Plymouth Rock (White) Brown Yellow Single Reddish bay Red Yellow White Primary use—Broiler.  New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; Black Used in production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in crossbreeding.	Mediterranean				P. 12				
Andalusian (Blue) White White Single Reddish bay White Dark slaty Blue Slaty blue used mainly for egg production beggn production of egg production of corossbreeding.  American Plymouth Rock (Barred) Brown Yellow Single Reddish bay Red Yellow Barred (sex-linked) Dual purpose; used in crosses sexing chicks a hatching.  Plymouth Rock (White) Brown Yellow Single Reddish bay Red Yellow White Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Red Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; bottom feet white Used in production of crossbreeding.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in crossbreeding.	Leghorn (white)	White	Yellow	Single	Reddish bay	White	Yellow	White	All three are small
American Plymouth Rock (Barred) Brown Yellow Single Reddish bay Red Yellow Barred (sex-linked) Dual purpose; used in crosses sexing chicks a hatching.  Plymouth Rock (White) Brown Yellow Single Reddish bay Red Yellow White Primary use—Broiler.  New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; bottom feet white crossbreeds.  Aslatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & black) crossbreeding.	Minorca (Black)	White	White	Single	Brown	White	Dark slate	Black	in size and are
Plymouth Rock (Barred)  Brown Yellow Single Reddish bay Red Yellow Barred (sex-linked)  Plymouth Rock (White)  Brown Yellow Single Reddish bay Red Yellow White Primary use— Broiler.  New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use— Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English  Cornish (White)  Brown Yellow Pea Pearl Red Yellow White Used in development or male lines for crossbreeding.  Australorp  Tinted White Single Brown Red Dark slate; bottom feet white production of crossbreeds.  Asiatic  Brahma (Light)  Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & black) crossbreeding.	Andalusian (Blue)	White	White	Single	Reddish bay	White	Dark slaty Blue	Slaty blue	used mainly for egg production— Leghorn is the mo popular.
(Barred) Brown Yellow Single Reddish bay Red Yellow Barred (sex-linked) Dual purpose; used in crosses sexing chicks a hatching.  Plymouth Rock (White) Brown Yellow Single Reddish bay Red Yellow White Primary use—Broiler.  New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; bottom feet white Used in production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in crossbreeding.									
Plymouth Rock (White) Brown Yellow Single Reddish bay Red Yellow White Primary use— Broiler. New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use— Broiler. Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; bottom feet white crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian (white & black) crossbreeding.									
(White)       Brown       Yellow       Single       Reddish bay       Red       Yellow       White       Primary use—Broiler.         New Hampshire       Brown       Yellow       Single       Reddish bay       Red       Yellow       Red       Primary use—Broiler.         Rhode Island Red       Brown       Yellow       Single       Reddish bay       Red       Yellow       Dark red       Dual purpose.         English       Cornish (White)       Brown       Yellow       Pearl       Red       Yellow       White       Used in development of male lines for crossbreeding.         Australorp       Tinted       White       Single       Brown       Red       Dark slate; bottom feet white       Black       Used in production of crossbreeds.         Asiatic       Brahma (Light)       Brown       Yellow       Pea       Reddish bay       Red       Yellow; Columbian (white & black)       Used in crossbreeding		Brown	Yellow	Single	Reddish bay	Red	Yellow	Barred (sex-linked)	used in crosses fo sexing chicks at
New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; bottom feet white Used in production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in crossbreeding.		Province	Vollous	Single	Paddish have	Pad	Vallous	White	Primanuso
New Hampshire Brown Yellow Single Reddish bay Red Yellow Red Primary use—Broiler.  Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; Black Used in production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in crossbreeding.	(AALIICE)	DIOWII	renow	Sirigle	neudisti Day	neu	TEHOW	AATIILE	
Rhode Island Red Brown Yellow Single Reddish bay Red Yellow Dark red Dual purpose.  English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; Black Used in production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in crossbreeding.	New Hampshire	Brown	Yellow	Single	Reddish bay	Red	Yellow	Red	
English Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; Black Used in bottom feet white  Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & black) Crossbreeding				29.0		\$1 <del>27127</del> .11	G-FESTIVE T	600077	
Cornish (White) Brown Yellow Pea Pearl Red Yellow White Used in development of male lines for crossbreeding.  Australorp Tinted White Single Brown Red Dark slate; Black Used in production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & black) crossbreeding.	Rhode Island Red	Brown	Yellow	Single	Reddish bay	Red	Yellow	Dark red	Dual purpose.
Australorp Tinted White Single Brown Red Dark slate; Black Used in production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & crossbreeding)	English								
Australorp Tinted White Single Brown Red Dark slate; Black Used in production of white crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & black) crossbreeding	Cornish (White)	Brown	Yellow	Pea	Pearl	Red	Yellow	White	development of male lines for
bottom feet production of crossbreeds.  Asiatic Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & black) crossbreeding	Australorp	Tinted	White	Single	Brown	Red	Dark slate:	Black	-
Brahma (Light) Brown Yellow Pea Reddish bay Red Yellow; Columbian Used in feathered (white & black) crossbreeding	The state of the s			Jingie		. 10.00	bottom feet		production of
feathered (white & black) crossbreeding	Asiatic								
	Brahma (Light)	Brown	Yellow	Pea	Reddish bay	Red			Used in crossbreeding for meat production
Source: U.S. Department of Agriculture. Agricultural Statistics, various years.									

#### C. Classification On the Basis of Production

On the basis of production chicken are of three types:

- 1) **Layer** is for egg production. Some popular layer breeds are Leghorn, Minorca, Ancona, Fayoumi, Isa Brown, Babycock, Star Cross, Lohman etc.
- 2) **Broiler** chicken are only for meat production. Plymouth rock, Cornish, Sussex, Dorking, Cochin, Brahma, Asil, Star Brow, Hi-line etc. are popular broiler breeds.

3) **Dual purpose (Egg and Meat) t**his types of breed are used for the purpose of both egg and meat production. Rhode Island Red, New Hampshire, Plymouth Rock etc. are popular breeds for both meat and egg production.

Most of the turkeys, ducks, and geese are raised for meat production. Except for hatching purposes, there is little market for turkey, duck, and goose eggs.

**Turkey breeds** including the Broad Breasted Bronze, Broad Breasted Large White, Beltsville Small White, Narragansett, Black, Slate, and Bourbon Red.

**Ducks breeds** the best breeds of ducks for meat production are the White Pekin, Aylesbury, Muscovy, Rouen, Cayuga, Swedish, and Call. The best egg-laying breeds are the Khaki Campbells and Indian Runners.

**Geese** the five common breeds of geese are the Toulouse, Embden, Chinese, Pilgrim, and African.